

# INDEX

## COPEIA, 1986, Nos. 1-4

### NEW NAMES

*Aguarunichthys torosus* 654-667  
*Amolops amoropalamus* 624-630  
*Amolops macrophthalmus* 623-629  
*Amolops orphnocnemis* 624-630  
*(Anchovieta)* 891, 895, 901  
*Apogon chalcus* 642-645  
*Bufo gargarizans popei* 572-577  
*Cyprinodon pachycephalus* 184-192  
*Etropus cyclosquamus* 142-150, 153, 154

*Leptobrotula breviventralis* 166-170  
*Notropis amecae* 170-183  
*Notropis aulidion* 170-183  
*Ololygon danae* 864-867  
*Ololygon exigna* 864, 866-869  
*Oryzias mekongensis* 473-478  
*Osornophryne talipes* 619-622  
*Pseudotropheus barlowi* 871-875  
*Similiparma* 858-862

### NEW AND ESTABLISHED NAMES

*Abies*  
*grandis* 398  
*sp.* 803  
*Abudefduf* 354, 857  
*luridus* 857  
*Acanthias vulgaris* 830  
*Acanthopis* 37  
*Acanthuridae* 277  
*Acanthurus* 355  
*Acara pulchra* 830  
*Acentromichthys* 654, 669  
*Acipenser*  
*guldenstadti persicus* 830  
*guldenstadti* × *nudiventris* 830  
*nudiventris* 830  
*sturio* 267  
*Acipenseridae* 830  
*Aclys* 526  
*Acosmanura* 260  
*Acris* 998, 1005  
*crepitans* 113, 995-999, 1005  
*gryllus* 1005  
*Acrochordidae* 424, 432  
*Acrochordoidea* 424  
*Acrochordus* 435, 918, 922-924  
*arafuræ* 238, 424-437  
*granulatus* 424, 432, 434, 435  
*javanicus* 427, 432-434, 916, 917, 922  
*Adioryx* 355  
*Aegires* 797  
*Aequidans pulcher* 830  
*Agalychnis dacnicolor* 113  
*Agama*  
*agama* 513  
*barbatus* 513  
*bibronii* 513  
*ornatus* 513  
*Agamidae* 33, 513  
*Aghistrodon*  
*contortrix* 432, 433, 807-811  
*contortrix contortrix* 808  
*contortrix laticinctus* 808-810

*contortrix laticinctus* × *contortrix pictigaster* 808  
*contortrix mokesen* 808, 810  
*contortrix pictigaster* 808-810  
*piscivorus* 432, 433  
*Aguarunichthys* 654, 657-659, 662, 663, 665, 667  
*torosus* 654, 659-664, 667  
*Algansea tincelli* 171, 177  
*Alligator mississippiensis* 116, 255, 333, 842  
*Allolobophora* 529  
*Allophryne* 262  
*Alnus* spp. 1002  
*Alopias vulpinus* 700-705  
*Alosa* 366  
*sapidissima* 683  
*Alutera scripta* 277-280  
*Alytes* 261, 837  
*muletensis* 837  
*Amazonsprattus* 899  
*Ambloplites constellatus* 392  
*Amblyglyphidodon* 356  
*Amblyopsidae* 1031  
*Amblyopsis spelaea* 1031  
*Amblyrhynchus cristatus* 66, 68  
*Ambrosia dumosa* 1011  
*Ambystoma* 28, 463, 464, 725, 726, 729  
*annulatum* 405  
*californiense* 263  
*gracile* 726  
*jeffersonianum* 404, 454-466  
*macrodictylum* 398, 405, 406  
*macrodictylum columbianum* 398-408  
*macrodictylum croceum* 405  
*macrodictylum sigallatum* 404, 405  
*maculatum* 404-406, 553, 729  
*opacum* 116, 404, 729  
*talpoideum* 404-406, 409  
*texanum* 405, 725-729  
*tigrinum* 405, 553, 726  
*Ambystomatidae* 398  
*Ameca splendens* 177, 830  
*Ameiva* 246, 530  
*ameiva* 451, 783, 804-806

- chrysolaema* 530  
 sp. 515  
*Amiurus nebulosus* 830  
*Ammodytes* 225  
   *americanus* 223  
   *hexapterus* 223–226  
   *marinus* 225  
   spp. 223  
*Ammodytidae* 360  
*Amniataba percoides* 427  
*Amolops* 623, 625, 629, 630  
   *afghanus* 630  
   *amoropalamus* 624–630  
   *cavilypanum* 623, 625, 630  
   *chunganensis* 630  
   *formosus* 630  
   *granulosus* 630  
   *hainanensis* 630  
   *jerboa* 623–630  
   *kinabaluensis* 623, 625, 630  
   *lifanensis* 630  
   *loloensis* 630  
   *macrophthalmus* 623–629  
   *orophnocnemis* 624–630  
   *phaeomerus* 623, 626–629  
   *poecilus* 623, 629  
   sp. 628  
   *whiteheadi* 623, 626–630  
*Amphibia* 550, 623  
*Amphibolurus*  
   *barbatus* 254, 513  
   *inermis* 513  
   *muricatus* 33  
   *reticulatus* 513  
   spp. 33  
*Amphiprion* 138  
   *chrysopterus* 135  
   *clarkii* 135–139  
   *melanopus* 135  
   spp. 138  
*Amphistichus argenteus* 312  
*Anabantidae* 830  
*Anabas*  
   *scandens* 830  
   *testudineus* 830  
*Anarbylus switaki* 263  
*Anarhichas* 711  
*Anas junius* 1005  
*Anchoa* 359, 891, 895, 898, 899  
   *argentivittata* 898  
   *belizensis* 898  
   *cayorum* 898  
   *chamensis* 898  
   *choerostoma* 898  
   *colonensis* 898, 899  
   *compressa* 898, 899  
   *cubana* 898  
   *curta* 898  
   *delicatissima* 898  
   *eigenmannia* 898  
   *elongata* 896–898  
   *exigua* 898  
   *filifera* 891, 895–901  
   *filifera filifera* 901  
   *filifera howelli* 901  
   *helleri* 898  
   *hepsetus* 898, 899  
   *howelli* 901  
   *ischana* 891, 893, 898  
   *januaria* 898  
   *lamprotaenia* 898, 899  
   *longipinna* 896–901  
   *lyolepis* 891, 896–901  
   *lyolepis lyolepis* 900  
   *lyolepis continentalis* 900  
   *lucida* 898  
   *marinii* 896–898  
   *mittchilli* 898  
   *mundeola* 898  
   *mundeoloides* 898  
   *naso* 891, 894–898  
   *nasus* 891, 894–898  
   *nasuta* 891, 895–900  
   *panamensis* 898  
   *parva* 898, 899  
   *platyargyrea* 901  
   *scofieldi* 898  
   *spinifer* 898  
   *starksi* 891, 893–901  
   *tricolor* 898  
   *trinitatis* 898, 899  
   *walkeri* 898  
*Anchovia* 895, 899  
   *macrolepidota* 893  
*Anchoviella* 898  
   *analisi* 898  
   *balboae* 898  
   *elongata* 901  
   *manamensis* 899  
*(Anchoviella)* 891, 895, 901  
*Ancylorhynchus* 359  
*Anguillidae* 513, 515  
*Anguilla*  
   *anguilla* 683  
   *bengalensis* 683, 684  
   *rostrata* 222, 366, 679–685  
*Anguillidae* 356, 359  
*Anguilliformes* 277  
*Anguis fragilis* 513  
*Anisochromidae* 711  
*Anniella*  
   *geronimensis* 263  
   *pulchra* 263, 513  
*Anolis* 76, 782, 963, 1042  
   cf. *armouri* 529  
   *brevirostris* 963  
   *carolinensis* 114, 451, 513, 961, 963  
   *chlorocyanus* 963–970  
   *coelestinus* 963  
   *cristatellus* 1042  
   *cybotes* 530, 963, 967  
   *distichus* 76–86, 530, 963, 965, 968, 970

- distichus dominicensis* 76–86  
*distichus floridanus* 76–86  
*ernestwilliamsi* 1042  
*sagrei* 84  
 spp. 698  
*stratulus* 513  
*Anomalopus* 37  
 spp. 33  
*Anoplarchus* 357  
 Antennariidae 218  
 Antennarioidae 218  
*Antigonia* 707  
*Antillophis parvifrons* 530  
*Anura* 503, 618, 623, 864  
*Apdemia* 797  
*Apeltes* 358, 677  
     *quadracus* 677  
 Aphredoderidae 1030  
*Aphredoderus* 1030  
     *sayanus* 1030, 1031  
*Apogon* 356, 641, 642  
     *apogonides* 643, 644  
     *chalcus* 642–645  
         *coccineus* 642  
         *evermanni* 641  
         *moluccensis* 643, 644  
         *monochrous* 643, 644  
         sp. 642  
 Apogonidae 354, 355, 359, 362, 641  
*Aporosaura anchietae* 514  
*Aprasia* 31, 36, 37, 526, 528  
     *parapulchella* 31, 36  
     *pulchella* 31, 32  
     *repens* 31, 32  
     *striolata* 31, 32  
 Archaeobatrachia 260  
*Arothron* 278  
     *hispidus* 277–280  
     *meleagris* 277–279  
*Arrhyton*  
     *exiguum* 530  
     *funereus* 530  
*Artedius* 357  
 Arthroleptidae 261  
*Ascaphus* 257  
     *truei* 438, 522  
*Astatoreochromis alluaudi* 312  
*Astatotilapia burtoni* 312, 321  
*Asterodactylus* 260  
*Astroedactylus* 261  
*Astronesthes* 495, 496  
*Astronotus*  
     *ocellaris* 312  
     *ocellatus* 830  
*Atelopus* 440–444, 618  
     *bufoniformis* 620, 621  
     *chiriquiensis* 443  
     *ebenoides* 622  
     *ignescens* 622  
     *oxyrhynchus* 443  
     *varius* 438–444  
*Atherina* 305  
     sp. 309  
 Atherinidae 359, 366  
 Atheriniformes 86, 227  
 Atherinomorpha 305  
*Atriplex spinifera* 1011  
*Aulopus* 707, 710  
     *japonicus* 707  
 Aulostomidae 277  
*Aulostomus chinensis* 277–280  
*Austrelaps superbus* 432  
  
*Balanus balanus* 541  
*Baleaphryne* 837  
 Balistidae 267  
*Barilus* 367  
*Barisia* 263  
*Barringtonia acutangula* 425  
*Bassobythites* 169  
*Bathygobius soporator* 213–216  
 Bathysauridae 361  
 Batrachoididae 831  
*Batrachoseps*  
     *aridus* 262  
     *attenuatus* 262  
     *campi* 262  
     *nigriventris* 262  
     *pacificus* ssp. 262  
     *simatus* 262  
     *stebbinsi* 262  
*Batrachylodes* 839  
*Bellator* 359  
*Belonesox belizanus* 89, 90, 301, 302  
 Belostomatidae 409  
*Benthoesema panamense* 548  
*Berberis* spp. 1002  
*Bergiaria* 654  
 Beryciformes 706, 707  
*Beryx* 707  
*Betta splendens* 540  
*Bitis* 1015, 1017, 1018  
     *arietans* 1015, 1017, 1018  
     *gabonica* 1015–1018  
     *nasicornis* 1015–1018  
 Blenniidae 207, 356  
 Blenniini 211  
*Blennius* 356  
*Boleichthys*  
     *whipplei* 637, 640  
     *whipplei montanus* 640  
*Bombinator* 260  
*Bombinator* 260  
*Borostomias* 495  
 Bothidae 140, 362  
 Brachionichthyidae 218  
*Brachydanio* 235  
     *rerio* 541  
 Brachydraconidae 363, 364  
*Brachyistius* 357  
*Brachyplatystoma* 654, 668, 670  
*Brevoortia* 359  
*Brooksia* 513

- Bufo* 408-423, 438  
*alvarius* 263  
*americanus* 25, 762  
*andrewsi* 562, 565, 567-577  
*arenarum* 113  
*bankorensis* 561, 563, 568-577  
*boreas* 398, 617  
*bufo andrewsi* 561  
*bufo asiaticus* 561  
*bufo bufo* 561-569  
*bufo gargarizans* 561  
*bufo minshanicus* 561  
*bufo wrightii* 562-569, 572-576  
*exsul* 263  
*gargarizans* 561, 562, 571-576  
*gargarizans gargarizans* 572, 573  
*gargarizans miyakonis* 561-576  
*gargarizans popei* 572-577  
*japonicus formosus* 561, 573, 576  
*japonicus japonicus* 561-576  
*marinus* 550  
*minshanicus* 562, 565-577  
*nelsoni* 263  
*obstetricans* 261  
*raddei* 571  
*regularis* 550  
*sachaliensis* 562-565, 571  
*terrestris* 408-423  
*tibetanus* 561-577  
*torrenticola* 561, 575, 576  
*vulgaris* 1008, 1009  
*woodhousei* 454-466  
*Bufonidae* 262, 439, 618  
*Bufonoidea* 262  
  
*Cacophis* 37, 38, 260  
*Cacopis* 260  
*Cacosternum boettgeri* 553  
*Calamita* 261  
*punctata* 260  
*Calamites* 261  
*Callager borneoensis* 750  
*Callanthias* 711  
*Callichthyidae* 830  
*Callichthys callichthys* 830  
*Callisaurus* 263  
*draconoides* 68, 513  
*Callopistes* 515  
*Calophysinae* 670, 671  
*Calophysus* 654, 657, 663, 665, 667, 668, 670  
*Calotes*  
*cristatellus* 513  
*versicolor* 513  
*Campostoma*  
*anomalum* 836  
*spp.* 392  
*Candoia* 839  
*Canis latrans* 822  
*Caproidae* 362  
*Capros* 707  
  
*Carangidae* 267, 277, 711  
*Caranx* 711  
*Carassius* 50, 235  
*auratus* 178, 830  
*Carcharhinidae* 829, 830  
*Carcharhinus*  
*leucas* 700  
*obscurus* 700-705  
*plumbeus* 700-705, 829-831  
*Carcharodon* 268  
*carcharias* 268, 700-705  
*Caretta* 56, 796, 798, 840  
*caretta* 53-57, 116, 247, 248, 750, 813  
*Carlia* 34, 37  
*gracilis* 33  
*novaeguineae* 33  
*spp.* 33  
*tetradactyla* 33  
*Carpophis* 431  
*vermis* 529  
*Carpiodes*  
*carpio* 227  
*sp.* 392  
*Catostomidae* 227, 364, 366  
*Catostomus*  
*commersoni* 836  
*occidentalis* 932  
*Cassiopeia* 797  
*xamachana* 796, 797  
*Caulophryniidae* 490, 491  
*Causus* 1015-1018  
*maculatus* 1015  
*Cebidichthys* 357  
*Centrarchidae* 227, 364, 366  
*Centrophryniidae* 490, 491  
*Centropomidae* 359, 579, 587  
*Centropominae* 579, 587  
*Centropomus* 579-611  
*affinis* 582, 607  
*altus* 582, 605  
*appendiculatus* 582, 594, 596  
*argenteus* 582, 594  
*armatus* 582-591, 604-606, 608  
*atridorsalis* 582, 606  
*brevis* 582, 606  
*constantinus* 582, 602, 603  
*cuvieri* 582, 591  
*ensiferus* 580, 582-591, 594, 605-609  
*gabbi* 582, 603  
*grandoculatus* 582, 591  
*heringi* 582, 600  
*macrochirus* 582, 607-609  
*medius* 582-593  
*mexicanus* 582-591, 601-605  
*nigrescens* 582-591, 596-598  
*parallelus* 582-591, 594, 601-605  
*pectinatus* 582-594  
*pedimacula* 582, 591  
*pellegrini* 582, 603  
*poeyi* 582-591, 599-602  
*robalito* 582-591, 606-608

- scaber* 582, 607  
*undecimalis* 581–591, 594–600  
*undecim-radiatus* 582, 594  
*unionensis* 582–591, 595–600  
*viridis* 582–591, 595–600  
*Centropyge* 712  
*tibicen* 136, 138  
*Centroscyrnus crepidater* 202–206  
*Ceratiadae* 480  
*Ceratias* 479–493  
*carunculatus* 488, 489  
*holboelli* 479–493  
*holboelli holboelli* 483, 484  
*holboelli sessilis* 485  
*holboelli tentaculatus* 482, 483, 492  
*(Paraceratias) mitsukurii* 488, 489  
*tentaculatus* 481–486, 491  
*uranoscopus* 481, 485–492  
*Ceratiidae* 479–481, 491  
*Ceratioidei* 480  
*Ceratioidei* 490  
*Ceratostethus bicornis* 308  
*Cereberus* 434, 918, 919, 922–926  
*rynclops* 916, 917, 919, 921–924  
*Cetengraulis* 899  
*juvensis* 899  
*mysticetus* 893  
*Cetorsorhamdia* 654, 667, 669  
*Chaenocephalus* 363  
*Chaetodontidae* 267, 355  
*Chalcides*  
*mionecton* 514  
*polylepis* 837  
*sexlineatus* 837  
*Chamaeleo*  
*basiliscus* 513  
*lateralis* 513  
*Chamaeleonidae* 513  
*Chamaelinorops* 530, 1042  
*Champscephalus* 363  
*Channa* 711  
*Channichthyidae* 363, 364  
*Characidae* 365  
*Characodon lateralis* 171, 178  
*Chauliodontidae* 361  
*Chauliodus* 495, 496, 499–501  
*Cheirocerus* 654, 661, 668–670  
*Chelodonia expansa* 1013  
*Chelonia* 56, 798  
*mydas* 53, 247, 254, 750, 797–802  
*mydas mydas* 247  
*Cheloniidae* 753  
*Chelydra* 741, 744, 746, 753–755  
*serpentina* 255, 741–756  
*Chilomeniscus*  
*cinctus* 263  
*punctatissimus* 263  
*savagei* 263  
*stramineus* 263  
*Chimaeridae* 360, 362  
*Chirostoma*  
*jordani* 177  
*mezquitali* 171, 178  
*Chirostomias* 495  
*Chondrichthyes* 202  
*Chriocentridae* 267  
*Chriope* 170  
*Chromis* 354, 356  
*Chrysemys* 263  
*picta* 698, 746, 749, 750, 782  
*Cichlasoma* 367  
*centrarchus* 216  
*citrinellum* 870, 874, 1028–1030  
*festivum* 830  
*managuense* 1030  
*nigrofasciatum* 830  
*Cichlidae* 311, 322, 365, 830, 870, 903  
*Cirrhitidae* 355  
*Citharichthys* 141, 359  
*arctifrons* 141, 150, 152–154  
*crossotus atlanticus* 151  
*micros* 149, 150  
*microstomus* 149  
*spilopterus* 150  
*unicornis* 150  
*Clarias garepinensis* 680  
*Claudius* 285, 288, 292  
*angustatus* 286–294  
*Clemmys*  
*guttata* 742–750  
*insculpta* 1001  
*Clinidae* 355  
*Clinocottus* 357  
*Clupea*  
*harengus* 25, 28, 714–719  
*harengus harengus* 714  
*harengus pallasi* 223, 714  
*pallasi* 714–719  
*Clupeidae* 267, 366  
*Cnemidophorus* 245, 246, 804, 806, 807  
*arubensis* 804, 806  
*dixonii* 263  
*flagellicaudis* 3, 263  
*gularis* 515  
*hypertythrus* 68  
*lemniscatus* 804–807  
*ocellifer* 783  
*sonorae* 263  
*sp.* 515  
*tigris* 3, 20, 68, 246, 515  
*Cobitidae* 830  
*Cobitis taenia* 830  
*Coelorhynchus* 362  
*Coleonyx* 263  
*sp.* 246  
*switaki* 263  
*variegatus* 3, 246, 513, 773, 781, 782  
*Colossoma macropomum* 193  
*Colostethus sp.* 622  
*Coluber* 812  
*constrictor* 432, 1001  
*mormon* 263

- Colubridae 432  
*Conger oceanicus* 222  
 Congridae 362  
*Conolophus* 67  
   *pallidus* 67, 68  
   *subcristatus* 66, 68  
*Cophosaurus* 263  
   *texanus* 3, 514  
*Coregonus* 128, 367  
   *lavaretus* 267  
*Corydoras palleatus* 830  
*Coryphaena hippurus* 276-281  
 Coryphaenidae 277  
*Coryphaenoides* 362  
*Coryphoblennius* 356  
 Cottidae 356-358  
*Cottus* 93, 365  
   *bairdi* 91-96  
   *caroliniae* 392, 394, 836  
   *gulosus* 380  
   *hypselurus* 380, 392  
   *ricei* 95  
*Cristiceps* 356  
*Crocodyl*† 843  
   *acutus* 747, 748  
   *johnstoni* 842  
   *niloticus* 332-341, 683  
   *siamensis* 332  
*Crotalus* 916, 919, 922, 925, 926  
   *atrox* 433  
   *catalinensis* 263  
   *cerastes* 433, 916, 917, 919, 922-926  
   *horridus* 433  
   *tortugensis* 263  
   *viridis* 433  
   *viridis oreganus* 816-820  
*Crotaphytus*  
   *bicinctores* 263  
   *collaris* 3, 513, 514, 980  
   *insularis* 263  
   *wislizenii* 263  
*Cryptoblepharus* spp. 33  
*Cryptobranchus alleganiensis* 553  
*Cryptopsaras* 479-482, 488-492  
   *couesi* 479, 481, 488, 489  
*Cryptosaras* 488  
*Cryptosarus* 488  
*Cryptosparas* 489  
   *atlantidis* 489  
   *normani* 489  
   *pennifer* 489  
   *valdiviae* 489  
*Ctenobrycon spilurus* 193  
*Ctenochaetus* 710  
*Ctenochatus* 355  
*Ctenopus* 37, 57, 60, 61, 450  
   *atlantis* 58, 60  
   *essingtonii* 33  
   *inornatus* 33  
   *lesueurii* 33  
   *regius* 58, 60, 61  
   *robustus* 33, 58, 60  
   spp. 33, 63  
   *taeniolatus* 58, 60, 61, 445-453  
   *uber* 58, 61  
*Cyanea* 797  
*Cyanothrissa* 367  
*Cyclopsetta* 141  
*Cycloramphini* 261  
*Cycloramphis* 261  
*Cyclorhamphini* 261  
*Cyclura* 67, 1042  
   *nubila* 65-69  
*Cymatogaster aggregata* 312  
 Cynoglossidae 362  
*Cynoscion* 359  
 (Cyprinella) 127  
*Cyprinidae* 219, 227, 359, 364-367, 830  
 Cypriniformes 227, 706  
*Cyprinodon* 45, 50, 51, 184, 189, 191  
   *bifasciatus* 190  
   *elegans* 189  
   *eximius* 184-192  
   *macrolepis* 190, 191  
   *macularius* 50  
   *maya* 190  
   *nevadensis* 50  
   *pachycephalus* 184-192  
 Cyprinodontidae 359, 367, 830  
 Cyprinodontiformes 706  
*Cyprinus carpio* 473, 677  
  
*Dactylethra* 261  
*Dactylethrinae* 261  
*Damalichthys* 358  
*Danio* 366  
*Darlingtonia* 529-531  
   *haetiana* 529-531  
*Dascyllus* 356  
*Deania calcea* 202-206  
*Decapterus* 712  
*Deirochelys reticularia* 782  
*Delma* 34, 36, 526, 528  
   *fraseri* 31, 32, 34, 36, 514  
   *inornata* 31, 32, 34, 36  
   *nasuta* 31, 32, 34, 36  
   *tincta* 528  
   *torquata* 528  
*Demansia* 37, 38  
   *psammophis* 33  
*Dendrobates* 260, 524  
   *pumilio* 438  
*Dendrochirus zebra* 136, 138, 686  
*Dermatemys* 292  
 Dermochelyidae 753  
*Dermochelys* 56  
   *coriacea* 53, 247-249, 796-798  
*Desmognathus* 236, 237, 731, 734, 736, 738, 739  
   *fuscus* 96, 235-237, 731-739  
   *monticola* 236, 237, 731-739  
   *ochrophaeus* 96, 236, 731-738  
   *quadramaculatus* 236, 731-739  
*Diadophis punctatus* 432, 529

- Diaphus* 362  
*Dicamptodon*  
   *aterrimus* 263  
   *copei* 263  
*Diceratiidae* 491  
*Diodon holocanthus* 277–281  
*Diodontidae* 277, 278  
*Diplodactylus* 526  
*Diplodus* 357  
*Diploglossus* 782  
   *lessonae* 782  
*Diplomystidae* 654  
*Diploprion* 711  
*Diporiphora*  
   *bilineata* 33  
   spp. 33  
*Dipsosaurus* 324–330  
   *dorsalis* 245, 514, 976  
*Discoglossidae* 261  
*Discoglossoidae* 260  
*Discoglossus* 837  
   *picta* 837  
*Ditrema temmincki* 312  
*Doderleinia* 707, 709, 711  
   *berycoidea* 706  
*Dormitator* 359  
*Dracaena* sp. 515  
*Duopalatinus* 654  
   *ollalae* 668  
*Dysticidae* 409
- Echidna zebra* 277–280  
*Echinorhinus cookei* 133  
*Echis* 1014, 1015, 1017  
   *carinatus* 1015  
*Egernia* 61  
   *cunninghami* 61, 445  
   *kingi* 3  
   *saxatilis* 61  
   *striolata* 58, 61, 63  
   *whitii* 58, 59, 61  
*Elaphe* 918, 921, 923, 925  
   *guttata* 432, 923  
   *guttata guttata* 924, 925  
   *obsoleta* 923  
   *obsoleta quadrivittata* 916, 917, 923–925  
   *vulpina* 812  
*Elapidae* 33, 264, 432  
*Eleotridae* 365  
*Eleotris* 359  
*Eleutherodactylus* 503, 662, 1042  
   *abbotti* 529, 530  
   *andi* 503  
   *armstrongi* 529, 530  
   *augusti* 263  
   *buckleyi* 622  
   *chloronotus* 622  
   *coqui* 520–523  
   *crassidigitus* 503–511  
   *cuaquero* 504–511  
   *curtipes* 621
- escoces* 505–507  
   *eunaster* 529, 530  
   *fitzingeri* 503  
   *glandulifer* 529, 530  
   *glaphycompus* 529  
   *heminota* 529  
   *longirostris* 504–511  
   *melanostictus* 504–511  
   *monnichorum* 504–511  
   *myersi* 621, 622  
   *oxyrhynchus* 529  
   *raniformis* 504–511  
   *rayo* 504–511  
   *rugulosus* 505–507  
   *sciagraphus* 529  
   sp. 621  
   *talamancae* 504–511  
   *vertebralis* 622  
   *whymperi* 622  
*Elgaria* 263  
   *multicarinata* 513  
*Elopidae* 359  
*Elosia nasuta* 261  
*Embiotoca* 311, 314, 317–320, 357, 358, 381  
   *jacksoni* 312–323, 381  
   *lateralis* 312–323, 381  
*Embiotocidae* 311, 322, 357, 358  
*Emmelichthyidae* 355  
*Enchelyopus cimbrius* 888  
*Engraulidae* 267  
*Engraulis* 899  
   *clarki* 891, 892, 895  
   *compressus* 898  
   *mordax* 40  
*Enhydrina schistosa* 432  
*Ephedra trifurca* 820  
*Epinephelinae* 580  
*Epinephelus* 709, 711  
*Epiplatys chaperi* 830  
*Eplatretus stoutii* 133  
*Eremias*  
   *lineocellata* 68  
   *lugubris* 68, 999  
   sp. 514  
*Eremiascincus* 59–63  
   *fasciolatus* 59–63  
   *richardsoni* 59–61  
   spp. 63  
*Eretmochelys imbricata* 53, 247, 248  
*Ericymba* 364  
*Erimyzon oblongus* 392, 836  
*Eristocophis* 1014–1017  
   *mcmahoni* 1014, 1015  
*Esocidae* 365, 367  
*Etheostoma* 365, 367, 637  
   *acuticeps* 538  
   *artesia* 631  
   *asprigene* 836  
   *blennioides* 392, 394, 538, 836  
   *caeruleum* 391, 392, 394, 538, 836  
   *camurum* 538  
   *chlorosomum* 836

- (Etheostoma)* 640  
*euzonum* 536  
*flabellare* 365, 835  
*gracile* 836  
*histrio* 836  
*juliae* 536-540  
*lepidum* 28  
*lynceum* 832-836  
*(Nanostoma)* 836  
*neopteron* 836  
*(Nothonotus)* 536, 538  
*(Oligocephalus)* 536, 631, 637  
*parvipinne* 836  
*podostemone* 365  
*pottsi* 178  
*proeliare* 544, 836  
*radiosum* 27, 631-641  
*radiosum cyanorum* 631  
*radiosum paludosum* 631  
*radiosum radiosum* 631  
*rufileineatum* 538  
*sp.* 545  
*spectabile* 18-30, 391, 392, 394, 538, 836  
*swaini* 836  
*tetrazonum* 536, 538  
*tippecanoe* 538  
*(Ulocentra)* 545, 836  
*whipplei* 631-641  
*whipplei alabamiae* 640  
*whipplei artesianae* 631-641  
*whipplei montanum* 631, 635  
*whipplei radiosum* 631  
*whipplei whipplei* 631-641  
*zonale* 392, 394, 832-836  
*Etmopterus baxteri* 202-206  
*Eutropus* 140-156, 359  
*crossotus* 140-156  
*cyclosquamus* 142-150, 153, 154  
*delsmani delsmanni* 141  
*delsmani pacificus* 141  
*ectenes* 141  
*intermedius* 141, 142  
*longimanus* 141, 150  
*micros* 150  
*microstomus* 140, 142, 143, 145, 146, 148-150, 153  
 154  
*peruvianus* 141  
*rimosus* 140, 142-150, 153, 154  
*Eumeces* 244, 245  
*fasciatus* 515  
*gilberti* 245  
*guttulatus* 512  
*kingi* 3  
*obsoletus* 512, 515  
*Eupomacentrus* 354  
*planifrons* 135  
*Eurycea* 97, 99, 102, 103, 107  
*bislineata* 97, 99, 102-104, 107  
*bislineata bislineata* 97, 99, 107  
*bislineata cirrigera* 97  
*bislineata rivicola* 97, 99, 107  
*bislineata wilderae* 97  
*n. sp.* 97, 99, 102-107  
*quadridigitata* 409  
*Euthynnus affinis* 42  
*Evermannichthys* 157-165  
*convictor* 159, 160  
*metzelaari* 157-165  
*silus* 159  
*spongicola* 157-165  
*Ferocactus peninsulæ* 518  
*Feylinia currori* 515  
*Ficimia* 263  
*Fistularia commersonii* 277-280  
*Fistularidae* 277  
*Flammeo* 355  
*Flourensia cernua* 820  
*Fundulus* 347, 359, 367, 709  
*catenatus* 392  
*grandis* 348, 349  
*heteroclitus* 342-351  
*olivaceus* 392, 394  
*spp.* 349  
*zebrinus* 87  
*Furina* 37, 38  
*Gadidae* 356-361, 886  
*Gadus* 360, 362  
*morhua* 888  
*Gaidropsarus* 356  
*mediterraneus* 826-829  
*Galaxias* 366  
*Galeocerdo cuvieri* 700-705  
*Galeomorphii* 202  
*Galeorhinus australis* 202-206  
*Gallotia* 837  
*Gallus gallus* 831  
*Gambelia* 263  
*wislizeni* 245, 514, 980  
*Gambusia* 471  
*affinis* 86, 89, 90, 227, 301, 392, 471, 534-536  
 937-945  
*alvarezii* 191  
*heterochir* 301  
*hurtadoi* 191  
*manni* 301  
*senilis* 191  
*yucatanana* 89  
*Gasterosteidae* 356, 359  
*Gasterosteus* 358, 365  
*aculeatus* 28, 946  
*volitans* 686  
*Gehyra* 526  
*mutilata* 781  
*oceanica* 781  
*spp.* 33  
*variegata* 33, 513  
*Gekko gekko* 513  
*Gekkonidae* 30, 33, 246, 513, 526-528



- Gekkoninae 961  
*Geochelone* 812  
*Gephyroberyx* 707  
*Gerres* 361  
*Gerrhonotus*  
   *cedrosensis* 263  
   *coeruleus* 263, 971  
   *liocephalus* 513  
   *multicarinatus* 263  
   *paucicarinatus* 263  
   spp. 263  
*Gigantactinidae* 490, 491  
*Gila* 1021, 1022  
   *bicolor* 49, 50  
   *bicolor mohavensis* 45-52  
   *cypha* 1021  
   *elegans* 1021  
   *orcutti* 45  
   *robusta* 1021  
   sp. 178  
*Gilchristella* 361  
*Girella* 709  
*Glossamia aprion* 427  
*Glyphidodon* 857, 858, 862  
   (*Parma*) 857  
   (*Parma*) *hermani* 857, 858  
*Glyphisodon* 857  
*Glyptophidium* 166, 169  
*Gnathanodon speciosus* 277-280  
*Gobiesocidae* 356  
*Gobiidae* 165, 213, 355-359  
*Gobiomorus* 359, 366  
*Gobiosoma* 159  
   *bosci* 213-216  
*Gobiosomini* 159  
*Gobius* 357  
   *juzo* 214, 215  
   *paganellus* 826  
*Goldiella* 654  
*Gonostomatidae* 361  
*Goodea cf. atripinnis* 177  
*Goodeidae* 830  
*Gopherus*  
   *agassizii* 263, 1010  
   *polyphemus* 787-795  
*Goslina* 654, 657, 668-670  
*Grammidae* 355, 711  
*Grammistidae* 711  
*Graptomys* 754  
*Grasseichthys* 848  
*Grayia* 434  
*Gulaphallus falcifer* 306  
*Gutierrezia* sp. 820  
*Gyalopion* 263  
*Gymnocephalus* 710  
*Gymnodactylus* 774, 775, 778-780  
   *geckoides* 773-786  
   *geckoides geckoides* 774  
*Gymnophthalmus* 783  
   *multiscutatus* 783  
   sp. 515  
*Gymnothorax pictus* 684  
*Gyrinidae* 409  
  
*Haemulon*  
   *flavolineatum* 269  
   *plumieri* 269  
*Hadrosauridae* 843  
*Halichoeres* 710, 712, 879  
*Halosauridae* 361  
*Haplochilus*  
   *chaperi* 830  
   *latipes* 830  
*Haplochromis* 367  
   *argens* 1029  
   *livingstoni* 830  
   *piceatus* 1029  
   *sauvagei* 1029  
*Harengula* 359  
*Harpagifer* 364  
*Harpagiferidae* 363, 364  
*Helisoma duyri* 541  
*Heloderma*  
   sp. 513  
   *suspectum* 512, 513, 515  
*Helodermatidae* 513  
*Helostoma* 710  
*Hemiaspis signata* 432  
*Hemibarbus* 709  
*Hemichromis bimaculatus* 216  
*Hemidactylus* 263, 774, 778-780  
   *flaviviridis* 513  
   *frenatus* 781  
   *mabouia* 773-786  
   *turcicus* 782, 956-962  
*Hemiergis* 61, 62  
   *decrensiensis* 58, 60-63, 445, 449  
   *peroni* 58, 60, 61, 445, 449, 450  
   *quadrilineatum* 60  
   spp. 59, 60  
*Hemisidae* 261  
*Hemisorubrum* 654, 668, 670  
*Hemistichodus* 848  
*Heptapterus* 654, 668-670  
*Heterandria* 86, 89, 90  
   *bimaculata* 88, 89  
   *formosa* 88-90, 301  
*Heterodon*  
   *nasicus* 432, 812  
   *platyrhinos* 432, 812  
*Heteronotia* 37, 840  
   *binoei* 33, 513  
*Heterophotus* 495  
*Hexagrammidae* 357  
*Hexagrammos stelleri* 224  
*Hexanchidae* 131, 362  
*Hexanchus*  
   *griseus* 131-135  
   *vitulus* 134  
*Hexanematchthys* 427, 433  
   *leptaspis* 427  
*Hilsa* 361

- Himantolophidae 490  
*Hippoglossoides* 707  
   *dubius* 711  
*Holbrookia* 263  
   *propinqua* 961, 980-986  
   *texana* 512  
 Holocentridae 355  
*Holocentrus* 355  
*Homalopsis buccata* 432  
*Hoplodactylus*  
   *maculatus* 971  
   *pacificus* 512, 513  
*Hybognathus* 367  
*Hybopsis*  
   *aestivalis* 227  
   *amblops* 392  
   *dissimilis* 392, 393  
*Hydrocotyle* 408-423  
   *umbellifera* 408-423  
*Hydrolagus collieri* 133, 205  
*Hydromantes* 263  
*Hydromantoides* 263  
 Hydrophiidae 264, 432  
*Hyla* 70-76, 802  
   *agilis* 864  
   *andersoni* 71-75, 454-466  
   *andersoni* × *cinerea* 70, 71, 75  
   *andersoni* × *femoralis* 70, 71  
   *arborea* 113  
   *avivoca* × *chrysoscelis* 70  
   *cadaverina* 263  
   *californiae* 263  
   *canastrensis* 864  
   *chrysoscelis* 532  
   *chrysoscelis* × *femoralis* 70  
   *cinerea* 70-75  
   *cinerea* × *gratiosa* 70  
   *crucifer* 109-115, 719-724  
   *cyanea* 261  
   *eximia* 263  
   *femoralis* 71-75  
   *fuscomarginata* 864  
   *gratiosa* 70, 73, 74  
   *maracaya* 864  
   *nebulosus* 864  
   *pseudopuma* 1007  
   *regilla* 398, 802-804, 930  
   *rosenbergi* 550  
   *rubra* 864  
   *smithii* 1004  
   *strigilata* 864  
   *versicolor* 113, 114, 532  
   *wrightorum* 263  
*Hylactophryne* 263  
*Hylaplesia* 260  
*Hylidae* 109, 864  
*Hymenocephalus* 362  
*Hynobius okiensis* 257  
*Hypentelium nigricans* 392, 395, 836  
*Hyperdon* 261  
*Hyperoliidae* 261  
*Hyperprosopon ellipticum* 312  
*Hypleurochilus geminatus* 207-213  
*Hypophthalminae* 670  
*Hypophthalmini* 670  
*Hypophthalmus* 654, 667, 669, 670  
*Hypsoblennius* 358  
   *hentzi* 207-213  
   *ionthas* 207-213  
*Hypsurus* 358  
*Hypsypops* 861, 862  
   *rubicunda* 135  
*Hysaplesia* 260  
  
*Ichthyomyzon gagei* 392  
*Ichthyophis* 257  
*Ictaluridae* 227, 364-367, 654, 830, 1024, 1027  
*Ictalurus*  
   (*Amiurus*) *melas* 1027  
   (*Amiurus*) *natalis* 1027  
   *dugesi* 177  
   (*Ictalurus*) *punctatus* 1027  
   *lupus* 646-652  
   *lupus* × *punctatus* 646-652  
   *natalis* 392, 540  
   *nebulosus* 540, 541  
   *punctatus* 227, 646-652  
   sp. 178  
*Iguana* 325, 328-330, 1042  
   *iguana* 116, 222, 324-332, 514  
*Iguanidae* 76, 245, 513, 516, 963  
*Iguanodon* 843  
*Iheringichthys* 654, 670  
*Ilyodon furcoides* 177  
*Imparfinis* 654, 667-669  
*Iniomi* 707  
*Ipnots murrayi* 271  
*Isurus oxyrinchus* 700-705  
  
*Katsuwonus pelamis* 42, 703  
*Kentropyx*  
   sp. 515  
   *striatus* 804-806  
*Kinosternon* 285, 288, 292, 293, 753  
   *acutum* 286-294  
   *alamosae* 271, 285-294  
   *angustipons* 285  
   *bauri* 285-294  
   *bauri bauri* 748, 750, 751  
   *creaseri* 285-294  
   *dunni* 285  
   *flavescens* 285-294, 812  
   *herrerai* 285-294  
   *hirtipes* 285-294  
   *integrum* 285-294  
   *leucostomum* 285-294  
   *oaxacae* 271, 285-294  
   *scorpioides* 285-294  
   *sonoriense* 285-294  
   *subrubrum* 285-294, 748-753  
*Klauberina* 263  
   *riversiana* 62

- Krameria* spp. 1011  
*Kyphosidae* 357
- Labeo* 847, 848  
*barbatus* 848  
*coubie* 847  
*cyclorhynchus* 847, 848  
*forkalii* 847  
*lividus* 848  
*macrostoma* 848  
*pseudocoubie* 847  
*variegatus* 847, 848
- Labidesthes* 367  
*sicculus* 392
- Labridae* 322, 355–358  
*Labroides* 322
- Lacerta* 837  
*agilis* 514, 838  
*lepida* 514, 838  
*perspicillata* 837, 838  
*schreiberi* 838  
*sp.* 514  
*spp.* 698  
*viridis* 514, 838  
*vivipara* 450, 451, 751
- Lacertidae* 514, 515
- Lacertilia* 30
- Lactoria diaphana* 136, 137
- Lampetra*  
*aepyptera* 392  
*appendix* 392  
*planeri* 205  
*tridentata* 133
- Lamprozelis*  
*calligaster* 432  
*getulis* 3, 807, 812  
*pyromelana* 3  
*triangulum* 812
- Lampropholis* 37  
*challengeri* 33  
*delicata* 33  
*guichenoti* 33  
*spp.* 33
- Lanius ludovicianus* 821
- Larrea tridentata* 1011
- Lasiognathus* 490, 491
- Lates* 427, 579, 580, 587  
*calcarifer* 427
- Laticaudidae* 432
- Latinae* 579, 587, 589
- Lavinia symmetricus* 932
- Lebistes reticulatus* 677, 946
- Leiarius* 667–670  
*marmoratus* 662, 668
- Leiognathidae* 267
- Leiopisma* 61  
*entrecaesteuxii* 58, 61  
*metallicum* 61  
*ocellatum* 61  
*trilineatum* 61
- Leiopelma* 839, 840
- Leiostomus* 359
- Lemmanura* 260
- Lepadogaster* 357
- Lepidochelys olivacea* 837
- Lepidopsetta bilineata* 224
- Lepomis* 366, 367  
*cyanellus* 392  
*humilis* 227, 229  
*macrochirus* 227, 392  
*megalotis* 189, 392  
*microlophus* 392
- Leptacanthichthys* 481
- Leptobrotula* 166–169  
*breviventralis* 166–170
- Leptodactylidae* 503
- Leptodeira* 530
- Leptophilypnus* 359
- Leptopis mexicanus* 432
- Lerista* 33, 37, 38  
*spp.* 33  
*stylus* 33
- Lethrinus* 709
- Leuciopimelodinae* 670, 671
- Leurognathus marmoratus* 734
- Lialis* 34–38, 526, 528  
*burtonis* 31–37, 514  
*jicari* 31, 32, 36
- Lichanura* 924  
*roseofuscus* 925
- Limnaeodius* 408–423  
*ocularis* 408–423
- Linophrynidae* 490, 491
- Liopsetta glacialis* 1024
- Liparidae* 356
- Lobulia*  
*morokana* 449  
*stanleyana* 449
- Lophiidae* 362
- Lophiiformes* 479
- Lophiophila* 261
- Lophiosilurus* 655
- Luciopimelodes* 655, 665, 667, 669  
*pai* 662, 664
- Lutjanidae* 267
- Lycengraulis* 898
- Lycium andersonii* 1011
- Lycodonomorphus* 433
- Lygodactylus* 774–780  
*klugei* 773–786
- Lygosoma taeniolum* 515
- Lysosominae* 61
- Mabuya* 782  
*heathi* 782
- Macrocephalus* 587
- Macroclernys temminckii* 116
- Macropodus* 235  
*opercularis* 830
- Macrostomias* 494, 500  
*furcatus* 503
- Macrouridae* 361, 362

- Makaira nigricans* 205  
*Malaclemys* 743, 744, 748, 754  
*terrapin* 116, 746-754  
*Malacosteus* 499  
*Manacopus falcifer* 306  
*Mancalias* 481, 483  
*bifilis* 482  
*kroyeri* 485  
*sessilis* 485  
*tentaculatus* 482  
*uranoscopus triflos* 485  
*xenistius* 485  
*Mancalius* 484  
*Markiana* (cf.) *nigripinnis* 193  
*Masticophis* 812, 924  
*aurigulus* 263  
*barbouri* 263  
*bilineatus* 3  
*flagellum* 432, 925  
*lateralis* 263  
*taeniatus* 116  
*Mauremys leprosa* 747  
*Megalaspis* 711  
*Megalonema* 655, 667-670  
*pauciradiatum* 670  
*platanum* 670  
*platycephalum* 669, 670  
*xanthum* 669, 670  
*Megalops cyprinoides* 427  
*Melanogrammus aeglefinus* 888, 946  
*Melanostigma atlanticum* 888  
*Melototaenia maculata* 427  
*Menticirrhus* 359  
*Merluccius productus* 133  
*Merlucciidae* 361  
*Merodonotus* 668, 670  
*Mesobatrachia* 260  
*Mesonanta insignis* 830  
*Microgadus tomcod* 888  
*Microglanis* 655  
*Micrometrus* 358  
*minimus* 312  
*Micropogonias* 359  
*Micropterus* 366, 367, 396  
*dolomieu* 392  
*punctulatus* 392  
*salmoides* 392  
*Microspathodon* 354  
*chrysurus* 857  
*frontatus* 862  
*minimus* 354  
*Micrurus* 1017  
*nigrocinctus* 1015, 1016  
*Miopsaras* 481  
*myops* 481, 487  
*Misgurnis fossilis* 830  
*Mochokidae* 830  
*Modiolus* 797  
*Moenkhausia dichroua* 193  
*Monacanthidae* 267, 277  
*Monomilopus* 169  
*Monopterus* 679  
*Morethia* 37  
*boulengeri* 33  
*Moridae* 361  
*Morone* 196-199, 709  
*americana* 196-199  
*chrysops* 196-199  
*saxatilis* 545  
*Moxostoma*  
*austrinum* 177  
*duquesnei* 392  
*erythrurum* 392, 836  
*Mugil*  
*cephalus* 227, 277-279, 343  
*curema* 277-279  
*Mugilidae* 227, 277, 359  
*Muraenidae* 277  
*Muraenolepidae* 363, 364  
*Mustela vison* 683  
*Mustelus*  
*canis* 829-832  
*lenticulatus* 202-205  
*Myctophiformes* 706, 707  
*Myoglanis* 655, 668, 670  
*Myopsaras* 481  
*Myosparas* 481  
*Myctophidae* 362  
*Myctophum* spp. 548  
*Myiobatrachus* 261  
*Mylossoma duriventris* 194  
*Myoxocephalus polyacanthocephalus* 224  
*Myripristis* 355, 356  
  
*Nandidae* 831  
*Nannorhamdia* 665, 667, 669  
*Nanostoma elegans* 836  
*Naso* 355  
*Natrix* 263  
*Naucrates* 711  
*Nectamia* 644  
*Nectophryniinae* 261  
*Necturus* 616  
*maculosus* 612-618, 762  
*Neetroplus* 367  
*Negaprion brevirostris* 700, 829-831  
*Nemachilus* 365  
*Nematistius* 712  
*Nematolosa* sp. 427  
*Neobatrachia* 260  
*Neobythitini* 169  
*Neoceratiidae* 490, 491  
*Neoditrema ransonetti* 312  
*Neogobius melanostomus* 213-216  
*Neonesithes* 495  
*Neoscopelus* 707  
*Neosilurus hyrtlii* 427  
*Neotrophus macrops* 150  
*Nerodia* 263, 433, 916, 918, 919, 921-926  
*cyclopion* 432  
*fasciata* 432, 754  
*fasciata clarki* 754

- fasciata compressicauda* 754  
*fasciata pictiventris* 916, 917, 919, 922-924  
*rhombifera* 432  
*sipidon* 241, 752  
*sipidon insularum* 757-772  
*sipidon sipidon* 757  
 sp. indet. 812  
*Nezumia* 362  
*Nippon* 579, 580, 711  
 (spinosus) 579  
*Nocomis*  
   *biguttatus* 392  
   *leptocephalus* 227  
*Notacanthidae* 362  
*Notechis* 839  
   *ater* 839  
*Notemigonus* 367  
   *crysoleucas* 227, 392  
*Notonectidae* 409  
*Notophthalmus* 464  
   *viridescens* 1006  
*Notorynchus cepedianus* 134  
*Notothenia* 363, 364  
*Nototheniidae* 363, 364  
*Notropis* 127-129, 170, 171, 177, 179, 235, 364-367  
   836  
   *aguirrequeuoi* 181  
   *altipinnis* 128  
   *amabilis* 227  
   *amecae* 170-183  
   *amnis* 128  
   *anogenus* 170  
   *ardens* 128  
   *aulidion* 170-183  
   *bellus* 128  
   *bifrenatus* 170, 171  
   *blennius* 170  
   *boops* 392, 836  
   *boucardi* 181  
   *buchanani* 171, 229  
   *calientis* 170-183  
   *calientis complex* 170-183  
   *camurus* 227, 835  
   *cayuga* 170  
   *chalybaeus* 128  
   *chihuahua* 191  
   *chrysocephalus* 392, 394  
   *chrysocephalus chrysocephalus* 836  
   *chrysocephalus isolepis* 835  
   *cornutus* 128  
   *cummingsae* 128  
   *emiliae* 128  
   *formosus* 171  
   *fumeus* 836  
   *galacturus* 392, 836  
   *garmani* 182  
   *greeni* 392  
   *heterodon* 170  
   *heterolepis* 128  
   *hubbsi* 171  
   *hypsipterus* 128  
   *imeldae* 181  
   *jordani* 170  
   *longirostris* 227  
   *lutrensis* 127, 227, 228  
   *lutrensis forlonensis* 182  
   (*Lythrurus*) 171  
   *maculata* 170  
   *moralesi* 181  
   *nazas* 181  
   *nubilus* 392  
   *ornatus* 181  
   *oxyrhynchus* 227  
   *ozarcanus* 392  
   *panarcys* 191  
   *pilsbryi* 392  
   *potteri* 227, 229  
   *procne* 128  
   *roseipinnis* 128  
   *rubellus* 392-394, 836  
   *sabinae* 392, 394  
   *sallei* 171, 181  
   *shumardi* 227, 229  
   *signipinnis* 129  
   *spilopterus* 122-130, 836  
   *spilopterus hypsisomatus* 122, 124  
   *spilopterus spilopterus* 122, 124  
   *stramineus* 128, 836  
   *telescopus* 392, 836  
   *tropicus* 181  
   *umbratilis* 128, 129  
   *umbratilis cyanocephalus* 836  
   *umbratilis umbratilis* 836  
   *venustus* 128, 227  
   *volucellus* 836  
   *welaka* 171  
   *whipplei* 127, 836  
*Noturus* 1027  
   *albater* 394  
   *exilis* 394  
   *gyrinus* 227, 229, 1027  
   *hildebrandi* 836  
   *miurus* 836  
   *munitus* 1027  
   *nocturnus* 836  
   *phaeus* 836  
   sp. 1027  
   *stigmatosus* 836  
*Odontaspis ferox* 166  
*Odontophrynus americanus* 532  
*Oedura*  
   *lessueurii* 513  
   *ocellata* 781, 961  
*Oligocottus* 357  
*Ololygon* 864, 867  
   *baumgardneri* 864, 867  
   *boesemani* 866, 867  
   *danae* 864-867  
   *egleri* 864  
   *exigua* 864, 866-869  
   *nebulosa* 864

- parkeri* 864, 866-869  
*rostrata* 866  
*rubra* 866  
*staufferi* 864, 867  
*trilineata* 864, 866  
*wandae* 867, 869  
*x-signata* 866  
*Oncorhynchus* 50  
*tshawytscha* 540  
*Oneirodidae* 490, 491  
*Onychomys* 821  
*leucogaster* 821  
*torridus* 821  
*Opheodrys* 431  
*aestivus* 432, 1001  
*Ophidiocephalus* 525, 528  
*taeniatus* 31  
*Ophisaurus koellikeri* 513  
*Opsanus*  
*beta* 831  
*tau* 910  
*Opsopoeodus emiliae* 836  
*Ornithopoda* 843  
*Oryzias* 473, 474, 477  
*celebensis* 473, 476, 477  
*curvinotus* 473, 476, 477  
*javanicus* 473, 476  
*latipes* 25, 309, 473, 476, 477  
*luzonensis* 473, 476, 477  
*mekongensis* 473-478  
*melastigma* 473, 476, 477  
*minutillus* 473, 476, 477  
*Osornophryne* 618, 621  
*bufoniformis* 618-622  
*percrassa* 618, 620  
*talipes* 619-622  
*Osteichthyes* 206  
*Osteopilus septentrionalis* 1007, 1008  
*Oxybelis*  
*aeneus* 432  
*fulgidus* 432  
*Oxyeleotris* 427, 433, 435  
*lineolatus* 427  
*Oxyjulis* 357  
*Oxylabrax* 587  
  
*Padogobius martensi* 213-216  
*Panaeus* spp. 680  
*Pandanus* 428  
*aquaticus* 425, 428  
*Pappogeomys merriami* 232  
*Parablennius* 356  
*marmoreus* 207-213  
*Parachaenichthys* 363  
*Paradelma* 526, 528  
*Paralabrax* 357  
*Paranthias* 711  
*Parapimelodus* 655, 670  
*Parastromateus* 712  
*Pariah* 159, 164  
*scotius* 157  
  
*Pariolius* 655, 679  
*Parma* 857-862  
*mccullochi* 861  
*microlepis* 861  
*occidentalis* 861  
*oligolepis* 861  
*polylepis* 861  
*unifasciata* 861  
*victoriae* 861  
*Parrichthys* 481  
*merrimani* 481, 487  
*Paulicia* 655, 668, 670  
*Pelamis* 434  
*platurus* 432, 434, 752  
*Pellonula* 367  
*Pelobatidae* 261  
*Pempheris* 267  
*Perca* 366, 707  
*flavescens* 43  
*loubina* 594  
*Percidae* 87, 227, 366, 367, 631  
*Perciformes* 227, 277, 706, 707  
*Percina* 365  
*caprodes* 28  
*macrolepidota* 87  
*maculata* 836  
*phoxocephala* 836  
*roanoka* 395  
*sciera* 28, 227, 836  
*vigil* 836  
*Percopsiformes* 1030  
*Percopsis* 366  
*Perrunichthys* 670  
*Perugia* 654, 667, 669  
*Petromyzon marinus* 205  
*Petropedetina* 261  
*Petrosaurus* 263  
*repens* 263  
*thalassinus* 263  
*Petrotilapia* 1029  
*Phallostethidae* 305  
*Phallostethus dunckeri* 308  
*Phanerodon* 357, 358  
*furcatus* 312  
*Phelsuma madagascariensis* 513  
*Phenacobius mirabilis* 836  
*Phenacostethus posthon* 308, 309  
*Philautus* 260  
*Philodryas*  
*nattereri* 432  
*olfersi* 432  
*Pholidae* 356  
*Pholis gunnellus* 222  
*Philoria frosti* 839  
*Photichthya* 495  
*Phoxinus* 365  
*erythrogaster* 392, 393, 836  
*Phratocephalus* 655  
*Phreatobius* 669  
*Phrynobatrachinae* 261  
*Phrynosoma*

- cerroense* 263  
*cornutum* 245, 514, 820–824  
*coronatum* 245  
*douglassi* 3, 245  
*mcalli* 245, 514  
*modestum* 820–824  
*orbiculare* 245  
*platyrhinos* 245  
*solare* 3  
 sp. 245  
*Phycis phycis* 826  
*Phyllobates* 524  
*Phyllodactylus* 263  
*geitje* 232  
*nocticolus* 263  
*porphyreus* 232  
*xanti* 263  
*Phyllopezus* 774, 775, 778–780  
*pollicaris* 773–786  
*pollicaris pollicaris* 774  
*Phyllurus* 526  
*Phrynobatrachus natalensis* 553  
*Physalia* 367  
*Physalaemus pustulosus* 265  
*Physalia* 797  
*Phython molurus* 265  
*Piaractus brachipomus* 194  
*Pimelodella* 655, 667  
*Pimelodidae* 365, 653  
*Pimelodina* 654–659, 661, 663, 665, 667, 668, 670  
*flavipinnis* 654–661, 663, 664  
*nasus* 654, 659–661  
*Pimelodinae* 671  
*Pimelodus* 655, 670  
*Pimphales* 366, 367  
*notatus* 227, 229, 392, 836  
*promelas* 392, 753, 755  
*Pinirampus* 654, 655, 667–670  
*agassizii* 669  
*barbancho* 669  
*pirinampu* 662, 664, 669  
*Pinus* 803  
*contorta* 802  
*jeffreyi* 803  
*ponderosa* 398  
*Pipa* 257, 260  
*Pipidae* 261  
*Pipoidea* 257  
*Pipoidei* 260  
*Pisces* 829, 903  
*Pituophis* 116, 120  
*melanoleucus* 3, 116–120, 812  
*melanoleucus melanoleucus* 116  
*Platichthys stellatus* 224  
*Platycephalidae* 86  
*Platynotus* 778  
*Platysilurus* 654, 668  
*Platystomatichthys* 654  
*Plectognathi* 707  
*Plectorhynchus pictus* 136, 138  
*Plectrypops* 355  
*Plesiopidae* 711  
*Plethodon* 96, 725, 728, 729  
*cinereus* 725–730  
*gordoni* 263  
*idahoensis* 263  
*stormi* 263  
*Pletholax* 526  
*gracilis* 31  
*Pleurodeles waltlii* 257  
*Pleuronectidae* 357, 359–361, 1024  
*Pleuronectiformes* 140, 706, 707  
*Podarcis*  
*bocagei* 837  
*muralis* 514  
*sicula* 837, 839  
*Poecilia* 90  
*butleri* 88, 89, 177  
*chica* 471  
*latipinna* 88–90, 295–304, 467–472, 534  
*reticulata* 88, 89, 301, 302, 471  
 sp. 88–90  
*sphenops* 471  
*Poeciliidae* 227, 359, 365, 534–536  
*Poeciliopsis*  
*gracilis* 301  
*infans* 171, 177  
*lucida* 301  
*monacha* 301  
*occidentalis* 301  
*prolifera* 301  
 sp. 88, 90  
*turneri* 301  
*Polycentrus schombbergkii* 831  
*Polychrus* 782  
*acutirostris* 782  
*Polyodon spathula* 205  
*Pomacanthidae* 136, 267  
*Pomacanthus* 712  
*Pomacentridae* 135, 311, 355–358  
*Pomacentrinae* 858  
*Pomacentrus* 356  
*wardi* 380  
*Pomadasyidae* 136, 138, 355, 358  
*Pomadasys*  
*crocro* 580  
*jubelini* 580  
*Pomoxis* 366  
*Priacanthidae* 277, 355  
*Priacanthus* 281  
*alalaua* 277, 279  
*arenatus* 277, 279  
*boops* 277, 279  
*cruentatus* 277–281  
 spp. 276  
*Priapella* 90  
*compressa* 88–90  
*Prieteila* 1026, 1027  
*phreatophila* 1024–1026  
*Prionace glauca* 202–206, 700–705  
*Prionotus* 359  
*Prionurus* 709

- Pristobrycon* 193-195  
*striolatus* 193-195  
*Prosopis juliflora* 820  
*Psammoperca* 579, 580, 587, 710  
*Psettodes* 707  
*Pseudacris* 408-423, 1008  
*ornata* 408-423  
*triseriata* 522  
*Pseudechis porphyriachus* 432  
*Pseudemoia spenceri* 61  
*Pseudemys* 263  
*nelsoni* 234, 235, 750  
*scripta* 691-700, 812  
*Pseudocercates* 1014  
*Pseudonaja* 839  
*nuchalis* 839  
*textilis* 432  
*Pseudophryne* 551  
*Pseudopimelodus* 654, 667, 670  
*Pseudoplatystoma* 654, 668, 670  
*Pseudopleuronectes americanus* 946  
*Pseudoraphis* 425  
*Pseudotropheus* 871, 873  
*aurora* 871  
*barlowi* 871-875  
*elegans* 871  
*greshakei* 871  
*livingstoni* 871  
*lombardoi* 871  
*zebra* 870, 871, 873  
*Pterengraulis* 898  
*atherinoides* 898  
*Pteroinae* 686  
*Pterois* 686, 689  
*cristatus* 686  
*miles* 686-690  
*muricata* 687  
*russelli* 689  
*volitans* 686-690  
*volitans castus* 686  
*zebra* 686  
*Pungitius* 358  
*Puntius* 366  
*Pygocentrus* 193-195  
*notatus* 193-195  
*Pygopodidae* 30-39, 514, 525-529  
*Pygopus* 32-36, 526-528  
*lepidopodus* 31-35, 526-528  
*nigriceps* 31-36, 526-528  
*Pyloodictis olivaris* 1027  
  
*Rachycentron canadum* 680  
*Raja* 359  
*clavata* 829, 830  
*Rajidae* 358, 363, 830  
*Ramphotyphlops* 37, 38, 839  
*Rana* 252, 408-423, 531, 551, 927, 932  
*areolata* 249, 252  
*arvalis* 249  
*aurora* 249, 252, 551, 927-931  
  
*aurora aurora* 928, 932, 933  
*aurora draytonii* 928-932  
*berlandieri* 263  
*blairi* 263  
*blythi* 524, 551  
*bombina* 260  
*boylii* 927-933  
*boylii boylii* 928  
*boylii muscosa* 928  
*boylii sierrae* 928  
*cascadae* 249, 252, 253, 927-933  
*catesbiana* 241, 522  
*chiricahuensis* 263  
*clamitans* 113, 114, 241  
*draytonii* 928  
*erythraea* 550  
*esculenta* 249  
*fisheri* 263  
*ibanorum* 524  
*kuhli* 524  
*microdisca* 524  
*microdisca finchi* 523-525  
*microdisca palavanensis* 523-525  
*muscosa* 927-933  
*onca* 263  
*palustris* 241, 531, 532  
*perezi* 837  
*pipa* 260  
*pipiens* 96, 114, 249, 252, 454, 931  
*pretiosa* 249, 252, 398, 551, 927-933  
*pretiosa pretiosa* 928  
*ridibunda* 837  
*septentrionalis* 239-244  
*sp.* 1006  
*sphenocephala* 249-253  
*sylvatica* 113, 241, 249-253, 454  
*temporaria* 249  
*tincloria* 260  
*utricularia* 408-423  
*virgatipes* 242  
*yavapaiensis* 262, 263  
*Raniceps raninus* 826, 828  
*Ranidae* 261  
*Ranidella* 840  
*Ranoidei* 260  
*Rasbora* 366  
*Reganichthys* 481  
*giganteus* 481, 484, 485  
*Reganula* 481  
*Regina* 431, 529  
*alleni* 431, 432  
*grahami* 432  
*rigida* 432  
*septemvittata* 432  
*Reptilia* 254  
*Rhacochilus* 357, 358  
*vacca* 312  
*Rhacophoridae* 261  
*Rhadinesthes* 495, 500  
*Rhamdia* 654, 670  
*Rheobatrachus*



- silus* 840  
*vittilinus* 840  
*Rhineura* 916, 917, 923, 924  
*floridana* 916, 917, 921  
*Rhinichthys*  
*atratus* 836  
*cataractae* 219–221  
*osculus* 380  
*Rhinophrynus* 257  
*Risor* 159, 164  
*Rutilus rutilus* 540  
  
*Sagittaria* 408–423  
*subulata* 408–423  
*Saiphos* 37  
*equalis* 33  
*Salmo* 365, 366  
*fario* 830  
*gairdneri* 946  
*trutta* 20, 25, 946  
*Salmonidae* 830  
*Salvadora hexalepis* 3  
*Salvelinus* 844  
*alpinis* 540, 945  
*fontinalis* 95, 945  
*malma* 945  
*namaycush* 945  
*Sarotherodon* 368  
*Sarpa* 357  
*Satan eurystomus* 1025  
*Sator* 245  
*angustus* 263  
*grandaevus* 263  
*Sauria* 963  
*Saurida* 876, 877, 883  
*longimanus* 876–882  
*sp. 1* 876, 879, 881  
*sp. 2* 876–883  
*spp.* 877, 881, 883, 884  
*undosquamis* 876–882  
*Sauromalus* 245, 519  
*australis* 263  
*hispidis* 518–520  
*obesus* 263, 514  
*sp.* 245  
*spp.* 263  
*Scaphiodontophis* 529  
*Scaphiopus* 263, 1009  
*bombifrons* 1008  
*couchi* 1008  
*holbrooki* 438, 1006  
*holbrooki hurteri* 1008  
*hurteri hurteri* 1008  
*multiplicatus* 263, 1008  
*Scaptochelys* 263  
*Scardinius erythrophthalmus* 540, 830  
*Scatophagus* 712  
*Sceloporus* 263  
*aeneus* 222  
*cyanogenys* 971, 977  
*graciosus* 68, 514, 698  
  
*grammicus* 514  
*jarrovi* 514, 515, 971–979  
*lineatus* 263  
*magister* 3, 263, 514  
*occidentalis* 68  
*olivaceus* 514  
*poinsetti* 514  
*sp.* 245  
*spinosus* 514  
*undulatus* 1–16, 514, 781, 1012–1014  
*virgatus* 68  
*zosteromus* 263  
*Sciaena undecimalis* 594  
*Sciaenidae* 355, 359  
*Scincidae* 33, 57, 245, 514, 515  
*Scincus officinalis* 515  
*Scomber japonicus* 39–43  
*Scomberoides* 711  
*Scophthalmus* 359  
*Scorpaena* 707  
*miles* 687  
*Scorpaenidae* 136, 138, 357, 358, 362  
*Scorpaeniformes* 706, 707  
*Scyliorhinus caniculus* 829, 830  
*Scyliorhynidae* 362, 830  
*Scyllium canicula* 829, 830  
*Sebastes* 357, 707  
*carnatus* 380  
*chrysomelas* 380  
*Semaprochilodus* 661  
*Semotilus atromaculatus* 392, 836  
*Serenoa repens* 1008  
*Seriola* 711  
*Serranidae* 355, 357, 358, 580, 711  
*Serranochromis robustus* 312  
*Serrasalmus* 193–195  
*rhombeus* 193–195  
*Sicyonia brevirostris* 680  
*Siganus* 709, 710  
*Siluriformes* 227  
*Simenchelyidae* 361  
*Similiparma* 858, 862  
*hermani* 858–863  
*Sistrurus catenatus* 433  
*Skiffia francescae* 177  
*Soleidae* 361  
*Solenophallus thessa* 305, 308, 309  
*Sonora*  
*bancroftae* 263  
*episcopa* 263, 432  
*mosaui* 263  
*semiannulata* 263  
*Sooglossidae* 262  
*Sorex* *sp.* 402  
*Sorubim* 654, 668–670  
*Sorubiminae* 670, 671  
*Sparidae* 359  
*Spartina alterniflora* 343, 743  
*Spea* 263  
*Spectrunculus grandis* 168  
*Sphaerodactylus* 530, 1042

- Sphenomorphus* 34, 37, 57, 61, 529  
     *crassicaudus* 33  
     *isolepis* 33  
     *kosciusko* 58, 60, 61  
     *quoyi* 58, 60, 61  
     *ruficaudus* 33  
     spp. 33  
*Sphyaena* 710  
     *aureoviridis* 594  
*Sphyrna lewini* 700-705  
*Sprattus sprattus* 94  
*Squalidae* 362, 830  
*Squalomorphii* 202  
*Squalus acanthias* 133, 202-206, 830, 993  
*Squatina*  
     *africana* 993  
     *californica* 987, 993  
     *dumeril* 993  
     *squatina* 992, 993  
*Staurois* 630  
*Staurotypinae* 288  
*Staurotypus* 285, 288, 292  
     *triporeatus* 286-294  
*Stenobranchius* 362  
*Sternoptychidae* 361  
*Sternotherus* 285, 288, 691, 753, 755  
     *carinatus* 285-294  
     *depressus* 285-294  
     *minor* 285-294  
     *odoratus* 285-294, 691, 698, 754-753, 812  
*Stilosoma extenuatum* 529  
*Stolephorus*  
     *naso* 894  
     *starksi* 892, 893  
*Stomatorhinus* sp. 825  
*Stomias* 494-503  
     *affinis* 494-503  
     *atriventer* 494-503  
     *boa* 494-503  
     *boa* 495  
     *boa ferox* 495  
     *brevibarbat* 494-503  
     *colubrinus* 495-503  
     *danae* 495-503  
     *gracilis* 495-503  
     *lampropeltis* 495-503  
     *longibarbat* 495-503  
     *nebulosus* 495-503  
     *pacificus* 495-503  
*Stomiidae* 494  
*Stomioidae* 494  
*Streptosaurus* 263  
     *tympanum* 58, 60, 61  
*Strongylura krefftii* 427  
*Syacium* 141  
*Synaphobranchidae* 361  
*Synbranchidae* 365  
*Synbranchus* 679  
*Syngnathidae* 355  
*Syngnathiformes* 277  
*Synodonhis nigrita* 830  
*Synodontidae* 876  
*Synodus englemani* 876  
*Tachydromus* sp. 514  
*Tantilla*  
     *hobartsmithii* 263  
     *relicta* 529  
     *yaquia* 263  
*Tapinurus semitaeniatus* 778  
*Tarentola*  
     *angustimentalis* 837  
     *boettgeri* 837  
     *gomerensis* 837  
     *mauritanica* 513  
*Taricha rivularis* 697  
*Tarletonbeania* 362  
*Tarpon atlanticus* 267  
*Tautoga onitis* 312  
*Tautoglabrus adspersus* 312, 321  
*Teiidae* 245, 515  
*Teleostei* 311, 870  
*Terapon* 709  
     *theraps* 709  
*Terrapene* spp. 1003  
*Tetraodontidae* 355  
*Tetraodontiformes* 277, 706, 707  
*Thamnophis* 437, 999  
     *butleri* 432  
     *digitatus* 263  
     *elegans* 402, 432, 999, 1001  
     *hammondi* 263  
     *marcianus* 999-1001  
     *radix* 812  
     *rufipunctatus* 432  
     *sauritius* 432  
     *sirtalis* 402, 432, 769, 807, 971  
     *sirtalis sirtalis* 916  
*Thaumatichthyidae* 490  
*Thryssa* 361  
*Thunnus*  
     *alalunga* 703  
     *albacares* 42  
     *mccoyi* 205  
     *thynnus* 703  
*Tilapia*  
     *aurea* 89, 177  
     (*Coptodon*) 905, 908  
     *mariae* 903-909  
     *mariae* × *zilli* 905  
     *melanotheron* 89  
     *nilotica* 312, 672-679  
     (*Pelmatolapia*) 904  
     *zilli* 903-909  
*Tilapia* 61  
     *cassurinae* 58  
     *nigrolutea* 61  
     *rugosa* 58-63, 515  
     *scincoides* 515  
     s. *scincoides* 515  
*Tinca*

- tinca* 830  
*vulgaris* 830  
 Tornieriobatinae 261  
*Trachemys* 751, 752  
*decussata* 748-751  
*Trachinotus* 359, 711, 712  
*Trachinus* 709  
*Trachyrhynchus* 362  
*Tramea* 1005  
*Trematomus* 363, 364  
*Triacanthodes* 707, 710  
 Trichuridae 362  
 Triglidae 362  
*Trigonolampa* 495  
 Trimorphodon  
   *biscutatus* 263  
   *lambda* 263  
   *lyrophanes* 263  
   *vandenburghi* 263  
   *vilkinsoni* 263  
*Trionyx* 753, 755  
   *muticus* 751, 753  
   sp. 752  
   *spiniferus* 751  
 Triportheus  
   *angulatus* 194  
   *elongatus* 194  
   *rotundatus* 194  
 Tripterygion 357  
 Trisopterus 360  
 Triturus  
   *helveticus* 404  
   *vulgaris* 404  
*Trogloglanis pattersoni* 1025  
*Tropidonotus natrix* 926  
*Tropidurus*  
   *delanonis* 984  
   *semitaeniatus* 783  
   *torquatus* 783  
*Tsuga canadensis* 1002  
*Tupinambis* sp. 515  
*Typhlina bramina* 839  
*Typhlobagrus* 654  
   *shufeldti* 481, 485  
*Typhloceratis* 481  
   *firthi* 481, 487  
*Typhlopsarus* 481  
*Typhlosparas* 481  
*Typhlopsaras* 481  
  
*Uma* 263  
   *inornata* 514  
 Umbridae 365, 367  
*Underwoodisaurus* 526  
*Unechis dwyeri* 432  
*Uperodon* 261  
*Urocyon cinereoargenteus* 822  
*Uromacer*  
   *catesbyi* 432  
   *oxyrhynchus* 432  
 Uromastix  
   *acanthinurus* 519  
   *hardwickii* 513  
*Urophycis* 380  
   *chuss* 380  
   *tenuis* 380  
*Urosaurus* 263  
   *microscutatus* 263  
   *nigricaudus* 263  
   *ornatus* 2, 3, 20, 514, 781  
*Uta* 263  
   *stansburiana* 1, 3, 514  
  
 Varanidae 515  
*Varanus* 330, 512  
   *gouldii* 515  
   *griseus* 515  
   *indicus* 515  
   *monitor* 515  
   *salvator* 330, 515  
   sp. 515  
   *varius* 67  
*Velella* 797  
*Vipera* 1014  
   *berus* 432, 837  
   *secaei* 837  
 Viperidae 432  
  
 Xantusia 263  
   *henshawi* 515  
   *vigilis* 3, 515  
 Xantusiidae 515  
 Xenoanura 260  
 Xenopinae 261  
 Xenopodinae 261  
*Xenopus* 257, 261  
   *laevis* 113, 263, 464, 617  
*Xenotoca melanosoma* 177  
*Xerepes* 357  
*Xiphister* 357  
*Xiphophorus*  
   *helleri* 89, 301, 471  
   *maculatus* 177, 301  
   *variatus* 301  
  
*Yuriria* n. sp. 177  
  
*Zanclus* 710  
   *canescens* 708  
   *cornutus* 276-279  
*Zebrasoma* 355  
*Zeiformes* 706, 707  
*Zenopsis* 707  
*Zeus* 707  
*Zoarcidae* 356, 357, 361  
*Zoogoneticus quitzeensis* 171, 177

# SUBJECT INDEX

## COPEIA, 1986, Nos. 1-4

**ACTIVITY**, *Sceloporus (undulatus, clarki)* (season. activ.) 1-18; *Cyclura nubila* (diel cycles relat. to thermoreg.) 65-69; *Ammodytes hexapterus* (pred. pressure varies with diel act. cycle) 223-226; *Sauromalus hispidus* (reduced by high extracell. K+) 518-520; *Necturus maculosus* (diel & season. cycles relat. to thermoreg.) 612-618; *Nerodia sipedon insularum* (seasonal.) 757-772.

**AGE**, *Sceloporus (undulatus, clarki)* (at matur.) 1-18; *Amphiprion clarkii*, *Centropyge tibicen*, *Lactoria diaphana*, *Dendrochirus zebra*, *Plectorhynchus pictus* (longevity of individuals at specific sites) 135-139; *Crocodylus niloticus* (age determ. by osteoderm laminae) 332-341; *Chelonia mydas* (at sex. matur.) 798-802; *Saurida (undosquamis, longimanus, sp. 1, sp. 2)* at sex. mat., short lifetime) 876-885; *Hemidactylus turcicus* (at sex. matur.) 956-962.

**BEHAVIOR**, *Padogobius martensi* (sounds made during courtship & terr. defense) 213-216; *Pseudemys nelsoni* (feeding rates higher in a group) 234-235; *Desmognathus fuscus* (substrate choice experiment) 235-237; *Rana utricularia*, *Limnaea ocularis*, *Pseudacris ornata*, *Bufo terrestris* (interspecific spatial avoidance of larvae) 408-423; *Atelopus varius* (homing & terr. after exp. manip. in natural hab.) 438-444; *Poecilia latipinna* (detection of fertile females by males) 467-472; *Sauromalus hispidus* (high extracell. K+ affects thermoreg. & feeding) 518-520; *Ictalurus nebulosus* (aggression suppressing pheromone) 540-544; *Necturus maculosus* (beh. thermoreg.) 612-618; *Anguilla rostrata* (anal. of spinning while feeding) 679-685; *Hyla crucifer* (adapt. signif. of breeding non-calling males) 719-724; *Ambystoma texanum* (absence of territor.) 725-730; *Chelydra serpentina*, *Malaclemys terrapin*, *Kinosternon subrubrum*, *Sternotherus odoratus*, *Clemmys guttata*, *Trionyx (muticus, spiniferus)*, *Chrysemys picta* (beh. osmoreg. in estuarine water) 741-756; *Crotalus viridis oreganus* (strike description) 816-820; *Mustelus canis* (tonic immobility) 829-832; *Pseudotropheus barlowi* n. sp. (gold color & territor.) 870-875; *Anolis chlorocyanus* (head bob displays) 963-970; *Sceloporus jarrovi* (thermoreg. beh.) 971-979; *Holbrookia propinqua* (male resp. to female colors) 980-986; *Clemmys insculpta* (stomping for earthworms) 1001-1004; *Osteopilus septentrionalis* (cannibalism) 1007-1009.

**BLOOD**, *Carcharodon carcharias*, *Isurus oxyrinchus*, *Carcharhinus plumbeus*, *obscurus*, *Galeocerdo cuvieri*, *Prionace glauca*, *Sphyrna lewini*, *Alpias vulpinus* (hemoglobin & hematocrit levels) 700-705.

**CALLS**, *Hyla (andersonii × cinerea, andersonii × femoralis)* (hybrid calls) 70-76; *Hyla crucifer* (calling vs non-calling males) 719-724; *Olophryotrogon (danae n. sp., exigua n. sp.)* 864-870; *Opsanus tau* (physiol. of sonic muscle) 910-915; *Rana (a. aurora, a. draytonii, boylei, cascadae, muscosa, pretiosa)* (anat. of vocal sacs) 927-936.

**CANNIBALISM**, *Osteopilus septentrionalis* (by larvae during metamorph.) 1007-1009.

**CHROMOSOMES**, *Hyla (andersonii × cinerea, andersonii × femoralis)* (kar. anal. of hybrids) 70-76; *Hybopsis aestivalis*, *Nocomis leptoccephalus*, *Notemigonus crysoleucas*, *Notropis (amabilis, buechanani, camurus, longirostris, lutrensis, oxyrhynchus, potteri, shumardi, venustus)*, *Pimephales notatus*, *Carpiodes carpio*, *Ictalurus punctatus*, *Noturus gyrinus*, *Gambusia affinis*, *Lepomis (humilis, macrochirus)*, *Percina sciera*, *Mugil cephalus* (GC-specific fluorochromes stain fish NORs) 226-231; *Oryzias mekongensis* (kar.) 473-478; *Rana palustris* (triploid individual) 531-533; *Gaidropsarus mediterraneus* (kar. var.) 826-829; *Prietella phreatophila* (kar.) 1024-1028.

**COLORATION**, *Hyla crucifer* (change to match background) 109-115; *Pseudotropheus barlowi* n. sp. (gold color & territor.) 870-875; *Holbrookia propinqua* (male resp. to female colors) 980-986; *Acris crepitans* (polymorph.) 995-999; *Hyla smithii* (of larval tail, relat. to pred.) 1004-1006.

**COMPETITION**, *Desmognathus (quadramaculatus, monticola, fuscus, ochrophaeus)* (for burrows, affects community compos.) 731-741.

**DENTITION**, *Hypsoblennius (ionthas, hentzi)*, *Hypleurochilus geminatus*, *Parablennius marmoratus* (relat. to diet) 207-213.

**DEVELOPMENT**, *Ambystoma jeffersonianum*, *Bufo woodhousei*, *Rana sylvatica*, *Hyla andersoni* (effect of low pH on embryos & larvae) 454-466; *Squatula californica* (embryonic) 987-994; *Cichlasoma citrinellum* (ontogen. changes in feeding morph.) 1028-1030.

**DISTRIBUTION**, *Scomber japonicus* (relat. to temp. tolerance) 39-44; *Etropus (cyclosquamus n. sp., crossotus, microstomus, rimosus)* (season. var.) 140-156; *Evermannichthys (metzelaari, spongicola)* (of sponge-dwelling fishes) 157-165; *Rana septentrionalis* (correlation of environ. requirements for reprod. and s. limit of distr.) 239-244; *Echidna zebra*, *Aulostomus chinensis*, *Fistularia commersonii*, *Mugil (cephalus, curema)*, *Priacanthus (alalaua, arenatus, boops, cruentatus)*, *Coryphaena hippurus*, *Gnathodon speciosus*, *Zanclus cornutus*, *Alutera scripta*, *Arothron (hispidus, meleagris)*, *Diodon holocanthus* (dispersal or vicariance in trans-Pacific

- allopat. popns.) 275-284; *Ambystoma jeffersonianum*, *Bufo woodhousei*, *Rana sylvatica*, *Hyla andersoni* (effect of low pH on local distr.) 454-466; *Ceratias* (*tentaculatus*, *holboellii*, *uranoscopus*), *Cryptoparas covei* 479-493; *Centropomus* (*pectinatus*, *medius*, *undecimalis*, *nigrescens*, *viridis*, *poei*, *parallelus*, *mexicanus*, *unionensis*, *armatus*, *robalito*, *ensiferus*) 579-611; *Osornophryne* (*talipes* n. sp., *percrassa*, *bufoniformis*) 618-622; *Etheostoma* (*whipplei*, *radiosum*) 631-641; *Pimelodina flavipinnis*, *Aguarunichthys torosus* 653-672; *Pterois* (*volitans*, *miles*) 686-690; *Clupea* (*harengus*, *pallasi*) (evidence of sympatry) 714-719; *Nerodia sipedon insularum* (on Lake Erie islands) 757-772; *Pseudotropheus barlowi* n. sp. 870-875; *Phycis chesleri* (Grand Bank, Labrador Shelf) 886-891; *Rana* (*a. aurora*, *a. draytonii*) (of vocal sac conditions) 927-936.
- ECOLOGY**, *Pituophis melanoleucus* (nest sites) 116-121; *Etropus* (*cyclosquamis* n. sp., *crossatus*, *microsomus*, *rimosus*) (temp., depth) 140-156; *Desmognathus fuscus* (exper. evidence of behav. niche expansion) 235-237; (resource partitioning in fishes) 352-388; (stream fish community recovery after flooding) 388-397; *Acrochordus arafurae* (life hist. aspects of a low-energy specialist) 424-437; *Ambystoma jeffersonianum*, *Bufo woodhousei*, *Rana sylvatica*, *Hyla andersoni* (sensitivity of embryos to low pH in lab & field) 454-466; *Nerodia sipedon insularum* (5-yr mark & recapt. study) 757-772; *Cnemidophorus lemniscatus* (ecol. factors affecting popn. density) 804-807.
- EGGS**, *Sceloporus* (*undulatus*, *clarki*) (size, clutch mass) 1-18; *Etheostoma spectabile* (size: seasonal var., relat. to # eggs/clutch and hatchling fitness) 18-30; *Poecilia latipinna* (var. in yolk loading time affects interbrood interval) 295-304; *Ambystoma jeffersonianum*, *Bufo woodhousei*, *Rana sylvatica*, *Hyla andersoni* (sensitivity of embryos to low pH in lab & field) 454-466; *Eleutherodactylus coqui* (season. diff. in temp. affects hatching time) 520-523; *Hemidactylus turcicus* (eggs adapted for colonization) 956-962; *Sceloporus undulatus* (effect of cold on devel. & surviv.) 1012-1014; *Aphredoderus sayanus* (carried in branchial cavity) 1030-1031.
- ELECTROPHORESIS**, *Anolis* (*distichus dominicensis*, *d. floridanus*) (detects subspecific intergradation) 76-86; *Eurycea* (n. sp., *b. bislineata*, *b. wilderae*, *b. cirrigera*, *b. rivicola*) (detects cryptic species) 96-108; *Morone* (*chrysops* × *americana*) (detects hybrids) 196-199; *Petromyzon marinus* (allelic var. in diff. age classes) 199-202; *Mustelus lenticulatus*, *Galeorhinus australis*, *Squalus acanthias*, *Deania calcea*, *Centroscymnus crepidater*, *Etmopterus baxteri*, *Prionace glauca* (low genet. var. in sharks) 202-207; *Echidna zebra*, *Aulostomus chinensis*, *Fistularia commersonii*, *Mugil* (*cephalus*, *curema*), *Priacanthus* (*alalaua*, *arenatus*, *boops*, *cruentatus*), *Coryphaena hippurus*, *Gnathanodon speciosus*, *Zanclus cornutus*, *Alutera scripta*, *Arothron* (*hispidus*, *meleagris*), *Di-*
- odon holocanthus* (genet. distance between trans-Pacific allopat. popns.) 275-284; *Kinosternon* (*bauri*, *subrubrum*, *herrerai*, *hirtipes*, *sonoriense*, *flavescens*, *alamosae*, *integrum*, *scorpioides*, *oaxacae*, *acutum*, *leucostomum*), *Staurotyphlops triporcatus*, *Claudius angustatus*, *Sternotherus* (*odoratus*, *carinatus*, *depressus*, *minor*) (systematics of family) 285-294; *Eleutherodactylus* (*fitzingeri*, *melanostictus*, *rayo*, *crassidigitus*, *talamancae*) (construct cladogram, PAUP program) 503-511; *Ictalurus* (*lupus*, *punctatus*) (evid. for valid sp.) 646-652; *Pseudemys scripta* (genet. diff. between popns.) 691-700; *Clupea* (*harengus*, *pallasi*) (elev. to sp. status) 714-719; *Gambusia affinis* (gen. structure of popns. in altered hab.) 937-945; *Acris crepitans* (compar. of color morphs) 995-999; *Eristocophis* (*mcMahonii*, *arietans*, *nasicornis*, *gabonica*), *Causus maculatus*, *Echis carinatus*, *Micrurus nigrocinctus* (construct cladogram, PAUP program) 1014-1020.
- ENDANGERED SPECIES**, *Gila elegans* (capture of individual, hypoth. for popn. decline) 1021-1023.
- EVOLUTION**, *Etheostoma spectabile* (possible factors selecting for var. egg sizes) 18-30; *Aprasia* (*parapulchella*, *pulchella*, *repens*, *striolata*), *Delma* (*fraseri*, *inornata*, *nasuta*), *Lialis* (*burtonis*, *jicari*), *Pygopus* (*lepidopodus*, *nigriceps*) (converg. evol. with snakes) 30-39; *Amphiprion clarkii* (possible selective pressures for protandry) 135-139; *Evermannichthys* (*metzelaari*, *spongicola*) (isolating mech. in geol. hist.) 157-165; *Rhinichthys cataractae* (morph. diff. between popns. isol. only 40 yrs) 219-221; *Desmognathus fuscus* (behav. niche expansion correlates with increased genet. var.) 235-237; *Poecilia latipinna* (fert. advertisement by females affects sexual selection) 467-472; *Pygopodidae* (selective factors in body elongation and limblessness) 525-529; *Tilapia nilotica* (est. of effects of selection on some heritable traits) 672-679; (evol. signif. of jaw muscle modif. in higher teleosts) 705-713; *Chelydra serpentina*, *Malaclemys terrapin*, *Kinosternon subrubrum*, *Sternotherus odoratus*, *Clemmys guttata*, *Trionyx* (*muticus*, *spiniferus*), *Chrysemys picta* (early stage of marine adap. in reptiles) 741-756; *Lygodactylus klugei*, *Phyllorhynchus pollicaris*, *Gymnodactylus gekoides*, *Hemidactylus mabouia* (evol. effects of invariable clutch size on life hist. charc.) 773-786; *Pseudotropheus barlowi* n. sp. (evol. of aggressively territorial gold cichlids) 870-875; *Anolis chlorocyanus* (of head bob displays) 963-970; *Holbrookia propinqua* (selective pressure for bright color in female iguanas) 980-986; *Eristocophis* (*mcMahonii*, *arietans*, *nasicornis*, *gabonica*), *Causus maculatus*, *Echis carinatus*, *Micrurus nigrocinctus* (evol. relationships) 1014-1020.
- FECUNDITY**, *Etheostoma spectabile* (relat. to egg size) 18-30; *Amphiprion clarkii* (lifetime) 135-139; *Fundulus heteroclitus* (size-specific var.) 342-351; *Squatina californica* 987-994.
- FEEDING**, *Caretta caretta* (relat. to hatchling swim-

- ming techniques) 53-57; *Tiliqua rugosa* (effect on preferred body temp.) 57-64; *Pseudemys nelsoni* (rates greater in a group) 234-235; *Acrochordus arafurae* (in a low-energy specialist) 424-437; *Sauromatus hispidus* (high extracell. K<sup>+</sup> reduces feeding in lab & field) 518-520; *Anguilla rostrata* (spinning to tear prey) 679-685; *Nerodia sipedon insularum* (seasonal., diet) 757-772; *Hyla regilla* (selection of pine pollen by larvae) 802-804; *Phycis chesteri* (diet & locale) 886-891; *Thamnophis marcianus* (large meal affects locomotion) 999-1001; *Clemmys insculpta* (stomping for earthworms) 1001-1004; *Cichlasoma citrinellum* (morph. of feeding apparat. changes during devel.) 1028-1030.
- FOOD, *Etheostoma spectabile*** (effects on fitness of hatchlings from var. egg sizes) 18-30; *Aprasia* (*parapulchella*, *pulchella*, *repens*, *striolata*), *Delma* (*fraseri*, *inornata*, *nasuta*), *Lialis* (*burtonis*, *jacari*), *Pygopus* (*lepidopodus*, *nigriceps*) (compar. of diets) 30-39; *Hexanchus griseus* (stomach contents) 131-135; *Pygocentrus notatus*, *Pristigaster striolatus*, *Serrasalms rhombeus* (changes during develop.) 193-196; *Darlingtonia haetiana* (extreme stenophagy) 529-531; *Dermochelys coriacea* (food requirements) 796-798.
- GENETICS, *Petromyzon marinus*** (allelic var. in diff. age classes) 199-202; *Mustelus lenticulatus*, *Galeorhinus australis*, *Squalus acanthias*, *Deania calcea*, *Centroscymnus crepidater*, *Etmopterus baxteri*, *Prionace glauca* (low genet. var. in sharks) 202-207; *Tilapia nilotica* (heritability estimates of 19 traits) 672-679; *Agkistrodon contortrix* (mult. paternity) 807-811; *Gambusia affinis* (gen. struct. of popns.) 937-945.
- GEOGRAPHIC LOCALITIES**
- Alabama, *Notropis* (*s. spilopterus*, *s. hypsissomatus*) 122-130; *Etheostoma* (*cyclosquamis* n. sp., *crossotus*) 140-156; *Sternotherus* (*depressus*, *minor*, *odoratus*) 285-294; *Etheostoma whipplei* 631-641.
- Alaska, *Ammodytes hexapterus* 223-226; *Rana pretiosa* 927-936; *Liopsetta glacialis* 1024.
- Arctic Ocean, *Clupea* (*harengus*, *pallasi*) 714-719.
- Arizona, *Sceloporus* (*undulatus*, *clarki*) 1-18; *Kinosternon* (*flavescens*, *sonoriense*) 285-294; *Phrynosoma* (*cornutum*, *modestum*) 820-824; *Sceloporus jarrovi* 971-979.
- Arkansas, (stream fish community) 388-397; *Etheostoma* (*whipplei*, *radiosum*) 631-641; *Etheostoma* (*lynceum*, *zonale*) 832-836.
- Ascension Is., *Chelonia mydas* 798-802.
- Atlantic, *Ceratiidae* (*tentaculatus* [S. ocean], *holboellii*, *uranoscopus*), *Cryptosaras couesi* 479-493; *Stomias* 494-503; *Centropomus* (*pectinatus*, *undecimalis*, *poyi*, *parallelus*, *mexicanus*, *ensiferus*) (W.) 579-611; *Clupea harengus* (N.) 714-719; *Phycis chesteri* (Grand Bank, Labrador Shelf) 886-891.
- Australia, *Aprasia* (*parapulchella*, *pulchella*, *repens*, *striolata*), *Delma* (*fraseri*, *inornata*, *nasuta*), *Lialis* (*burtonis*, *jacari*), *Pygopus* (*lepidopodus*, *nigriceps*) 30-39; *Ctenotus* (*atlas*, *regius*, *robustus*, *uber*), *Egernia striolata*, *Eremiascincus* (*fasciolatus*, *richardsoni*), *Hemiergis* (*decreasensis*, *peroni*), *Tiliqua rugosa* (S.A.), *Ctenotus taeniolatus*, *Egernia whitii*, *Leiopisma entrecasteauxii*, *Sphenomorphus* (*kosciuskoii*, *quoyi*, *tympanum*), *Tiliqua cassurinae* (N.S.W.) 57-64; *Acrochordus arafurae* (N. Terr.) 238-239; *Acrochordus arafurae* (N. Terr.) 424-437; *Ctenotus taeniolatus* (N.S.W.) 445-453; *Pygopodidae* 525-529; *Sauvrida* (*undosquamis*, *longimanus*, sp. 1, sp. 2) (north-west shelf) 876-885.
- Belize, *Evermannichthys metzelaari* 157-165; *Cladius angustatus*, *Staurotyphlops triporcatus*, *Kinosternon* (*leucostomum*, *scorpioides*) 285-294.
- Borneo, *Rana microdiscus finchi* 523-525; *Amolops* (*amoropalamus* n. sp., *poecilus*, *whiteheadi*, *orphnomenis* n. sp., *phaeomerus*, *jerboa*, *macrophthalmus* n. sp.) 623-630.
- Brazil, *Pimelodina flavipinnis* 653-672; *Lygodactylus klugei*, *Phyllorhynchus pollicaris*, *Gymnodactylus geckoides*, *Hemidactylus mabouia* 773-786; *Cnemidophorus lemniscatus* 804-807.
- California, *Scomber japonicus* 39-44; *Gila bicolor mohavensis* 45-52; *Hexanchus griseus* 131-135; *Sauromatus* sp., *Gambusia wislizeni*, *Sceloporus* sp., *Phrynosoma* sp., *Eumeces gilbertii*, *Cnemidophorus*?, *Coleonyx* sp. (fossils) 244-246; *Mugil cephalus* 275-284; *Embiotoca* (*jacksoni*, *lateralis*) 311-323; *Hyla regilla* 802-804; *Rana* (*aurora*, *boylei*, *cascadeae*, *muscosa*) 927-936; *Squatina californica* 987-994; *Gopherus agassizii* 1010-1012.
- Canada, *Notropis* (*s. spilopterus*, *s. hypsissomatus*) (Ontario) 122-130; *Rana septentrionalis* 239-244; *Phycis chesteri* (Labrador) 886-891; *Rana* (*aurora*, *pretiosa*) (Brit. Columbia) 927-936; *Salvelinus alpinus* (Northwest Terr.) 945-955.
- Cape Verde Archipelago, *Simuliparma* n. gen. *hermani* 857-863.
- China, *Bufo* (*tibetanus*, *minshanicus*, *g. gargarizans*, *g. popei*, *andrewsi*) 561-579.
- Colombia, *Evermannichthys metzelaari* (Isla de Providencia) 157-165; *Eleutherodactylus* (*fitzingeri*, *longirostris*, *raniformis*, *crassidigitus*) 503-511; *Osornophryne* (*percrassa*, *bufoniformis*) 618-622; *Pimelodina flavipinnis* 653-672.
- Colorado, *Sceloporus undulatus* 1-18; *Gila elegans* 1021-1023.
- Connecticut, *Sternotherus odoratus* 285-294.
- Costa Rica, *Atelopus varius* 438-444; *Eleutherodactylus* (*andi*, *fitzingeri*, *cuaquero*, *melanostictus*, *rayo*, *crassidigitus*, *talamancae*) 503-511; *Chelonia mydas* 798-802.
- Cuba, *Cyclura nubilus* (introd. to a Puerto Rican island) 65-69.
- Cyprus, *Caretta caretta* 53-57.
- Easter Island, *Apogon chalcinus* n. sp. 641-645.
- Ecuador, *Eleutherodactylus longirostris* 503-511; *Osornophryne* (*italipes* n. sp., *bufoniformis*) 618-622.



- El Salvador, *Kinosternon scorioides* 285-294.
- Florida, *Hyla (andersonii) × cinerea, andersonii × femoralis* 70-76; *Anolis (distichus dominicensis, d. floridanus)* (colonization from Bahamas and Hispaniola) 76-86; *Etropus (cyclosquamus n. sp., crossotus, microstomus, rimosus)* 140-156; *Evermannichthys spongicola* 157-165; *Pseudemys nelsoni* 234-235; *Coryphaena hippurus* 275-284; *Sternotherus (minor, odoratus)*, *Kinosternon bauri* 285-294; *Poecilia latipinna* 295-304; *Rana utricularia*, *Limnaeus ocularis*, *Pseudacris ornata*, *Bufo terrestris* 408-423; *Poecilia latipinna* 467-472; *Gopherus polyphemus* 787-795; *Caretta caretta* 813-816; *Tilapia (zilli, mariae)* 903-909; *Osteopilus septentrionalis* 1007-1009.
- Galapagos Is., *Aulostomus chinensis*, *Zanclus cornutus*, *Arothron meleagris* 275-284.
- Georgia, *Sceloporus undulatus* 1-18; *Etropus (cyclosquamus n. sp., crossotus, microstomus, rimosus)* 140-156; *Fundulus heteroclitus* (Sapelo Is.) 342-351; *Anguilla rostrata* 679-685; *Gopherus polyphemus* 787-795.
- Grand Cayman Is., *Priacanthus cruentatus* 275-284.
- Guatemala, *Kinosternon leucostomum* 285-294.
- Guyana, *Kinosternon scorioides* 285-294.
- Hawaii, *Leptobrotula n. gen. breviventralis n. sp.* 166-170; *Echidna zebra*, *Aulostomus chinensis*, *Fistularia commersonii*, *Mugil cephalus*, *Priacanthus cruentatus*, *Coryphaena hippurus*, *Gnathodon speciosus*, *Zanclus cornutus*, *Alutera scripta*, *Arothron (hispidus, meleagris)*, *Diodon holocanthus* 275-284.
- Hispaniola, *Darlingtonia haetiana* 529-531; *Anolis chlorocyanus* 963-970.
- Honduras, *Kinosternon leucostomum* 285-294.
- Idaho, *Ambystoma macrodactylum columbianum* 398-408; *Rana pretiosa* 927-936.
- Illinois, *Notropis (s. spilopterus, s. hypsismatus)* 122-130; *Etheostoma (lynceum, zonale)* 832-836.
- Indian Ocean, *Ceratas (tentaculatus [S. ocean], holboellii, uranoscopus)*, *Cryptosaras couesi* 479-493; *Stomias* 494-503; *Pterois miles* 686-690.
- Indiana, *Eurycea n. sp.* 96-108; *Sternotherus odoratus* 285-294; *Gambusia affinis* 534-536.
- Iowa, *Etheostoma zonale* 832-836.
- Italy, *Padogobius martensi* 213-216; *Gaidropsarus mediterraneus* 826-829.
- Japan, *Amphiprion clarkii*, *Centropyge tibicen*, *Lactoria diaphana*, *Dendrochirus zebra*, *Plectorhynchus pictus* 135-139; *Bufo (j. japonicus, j. formosus, torrenticola, gargarizans miyakonis)* 561-579.
- Kansas, *Sceloporus undulatus* 1-18; *Etheostoma whipplei* 631-641; *Acris crepitans* 995-999.
- Kentucky, *Notropis (s. spilopterus, s. hypsismatus)* 122-130.
- Korea, *Bufo g. gargarizans* 561-579.
- Lake Malawi, *Pseudotropheus barlowi n. sp.* 870-875.
- Lake Michigan, *Cottus bairdi* 91-96.
- Louisiana, *Heterandria formosa* 86-91; *Sternotherus carinatus* 285-294; *Etheostoma whipplei* 631-641; *Ambystoma texanum* 725-730.
- Maine, *Conger oceanicus* 222-223; *Rana septentrionalis* 239-244.
- Maryland, *Notropis (s. spilopterus, s. hypsismatus)* 122-130; *Hyla crucifer* 719-724.
- Massachusetts, *Eurycea bislineata* 96-108; *Sternotherus odoratus* 285-294.
- Mexico, *Heterandria (bimaculata, jonesi)*, *Xiphophorus (maculatus, helleri)* (Veracruz), *Poeciliopsis sp.*, *Priapella compressa*, *Poecilia bulteri*, *Profundulus (labialis, punctatus)* (Chiapas), *Gambusia yucatana* (Quin. Roo), *Belonesox belizanus* (Tabasco) 86-91; *Hexanchus griseus* (Baja Cal.) 131-135; *Notropis amecae n. sp.* (Jalisco), *Notropis audidion n. sp.* (Durango), *Notropis calientis* 170-183; *Cyprinodon pachycephalus n. sp.* (Chihuahua) 184-192; *Sceloporus arneus* 232-233; *Echidna zebra*, *Fistularia commersonii*, *Mugil curema*, *Priacanthus cruentatus*, *Coryphaena hippurus*, *Gnathodon speciosus*, *Zanclus cornutus*, *Arothron meleagris*, *Diodon holocanthus* (Gulf of Cal.) 275-284; *Staurotyptus triporcatus* (Oaxaca), *Kinosternon acutum* (Veracruz), *K. alamosae* (Sonora), *K. flavescens* (Chihuahua), Durango, Sonora, Tamaulipas), *K. herrerae* (Veracruz), *K. hirtipes* (Chihuahua, Durango, Guanajuato, Jalisco, Michoacan), *K. integrum* (Colima, Jalisco, Michoacan, Oaxaca, Queretaro, Sinaloa, Sonora, Tamaulipas), *K. leucostomum* (Veracruz), *K. oaxacae* (Oaxaca), *K. scorioides* (Chiapas, Tamaulipas, Yucatan), *K. sonoriense* (Chihuahua) 285-294; *Ictalurus lupus* (Tamaulipas) 646-652; *Rana aurora* (Baja Cal.) 927-936; *Hyla smithii* (Jalisco) 1004-1006; *Prietella phreatophila* (Coahuila) 1024-1028.
- Michigan, *Hyla crucifer* 109-115; *Morone (chrysops × americana)* 196-199; *Petromyzon marinus* 199-202; *Rana septentrionalis* (U.P.) 239-244.
- Minnesota, *Notropis (s. spilopterus, s. hypsismatus)* 122-130; *Rana septentrionalis* 239-244; *Etheostoma zonale* 832-836.
- Mississippi, *Etropus (cyclosquamus n. sp., crossotus)* 140-156; *Sternotherus carinatus* 285-294; *Etheostoma whipplei* 631-641; *Etheostoma (lynceum, zonale)* 832-836.
- Missouri, *Notropis (s. spilopterus, s. hypsismatus)* 122-130; *Etheostoma juliae* 536-540; *Etheostoma whipplei* 631-641; *Etheostoma (lynceum, zonale)* 832-836.
- Montana, *Rana pretiosa* 927-936.
- Nebraska, *Sceloporus undulatus* 1-18; *Rhinichthys cataractae* 219-221; *Kinosternon flavescens* 285-294.
- Nevada, *Rana (aurora, pretiosa)* 927-936.
- New Hampshire, *Rana septentrionalis* 239-244.
- New Jersey, *Pituophis melanoleucus* 116-121; *Etropus microstomus* 140-156; *Kinosternon subrubrum* 285-294; *Bufo woodhousei*, *Hyla andersoni* 454-466; *Chelydra serpentina* 741-756.
- New Mexico, *Sceloporus undulatus* 1-18; *Kinosternon flavescens* 285-294.
- New York, *Notropis (s. spilopterus, s. hypsismatus)* 122-130; *Etropus microstomus* 140-156; *Rana sep-*

- tenriorialis* 239–244; *Sternotherus odoratus* 285–294; *Carcharodon carcharias*, *Isurus oxyrinchus*, *Carcharhinus plumbeus*, *obscurus*, *Galeocerdo cuvieri*, *Prionace glauca*, *Sphyrna lewini*, *Alopias vulpinus* 700–705.
- New Zealand, *Mustelus lenticulatus*, *Galeorhinus australis*, *Squalus acanthias*, *Deania calcea*, *Centroscymnus crepidater*, *Etmopterus baxteri*, *Prionace glauca* 202–207.
- Nicaragua, *Eleutherodactylus (fitzingeri, talamancae)* 503–511; *Cichlasoma citrinellum* 1028–1030.
- North Carolina, *Etmopterus (cyclosquamus n. sp., crossotus, microstomus, rimosus)* 140–156; *Hyposoblennius (ionthas, hentzi)*, *Hypleurochilus geminatus*, *Parablennius marmoratus* 207–213; *Desmognathus fuscus* 235–237; *Kinosternon subrubrum* 285–294; *Rana palustris* 531–533; *Desmognathus (quadramaculatus, monticola, fuscus, ochrophaeus)* 731–741; *Aphredoderus sayanus* 1030–1031.
- Ohio, *Sceloporus undulatus* 1–18; *Eurycea (n. sp., b. bislineata, b. wilderae, b. cirrigera, b. rivicola)* 96–108; *Notropis (s. spilopterus, s. hypsiosomatus)* 122–130; *Morone (chrysops × americana)* 196–199; *Nerodia sipedon insularum* 757–772; *Etheostoma (lynceum, zonale)* 832–836.
- Oklahoma, *Etheostoma (whipplei, radiosum)* 631–641; *Heterodon (nasicus, platyrhinus)*, *Coluber* or *Masticophis*, *Elaphe vulpina*, *Lampropeltis (getulus, triangulum)*, *Pituophis melanoleucus*, *Thamnophis cf. radix*, *Nerodia sp. indet. (fossils)* 811–812.
- Oregon, *Rana (aurora, boylei, cascadae, pretiosa)* 927–936.
- Pacific, *Ceratiis (tentaculatus [S. ocean], holboellii, uranoscopus)*, *Cryptopsaras coxeti* 479–493; *Stomias* 494–503; *Centropristis (medius, nigrescens, viridis, unionensis, armatus, robalito) (E.)* 579–611; *Pterois volitans* 686–690; *Clupea pallasi (N.)* 714–719.
- Panama, *Priacanthus cruentatus*, *Coryphaena hippurus*, *Alutera scripta*, *Arothron (hispidus, meleagris)* 275–284; *Kinosternon leucostomum* 285–294; *Eleutherodactylus (fitzingeri, longirostris, raniformis, melanostictus, monnichorum, crassidigitus)* 503–511.
- Pennsylvania, *Notropis (s. spilopterus, s. hypsiosomatus)* 122–130; *Ambystoma jeffersonianum*, *Rana sylvatica* 454–466; *Clemmys insculpta* 1001–1004.
- Peru, *Aguarunichthys torosus*, *Pimelodina flavipinnis* 653–672.
- Philippine Is., *Zanclus cornutus* 275–284; *Manacopus falcifer* 305–310.
- Puerto Rico, *Cyclura nubila* (introd. to Isla Magüeyes) 65–69; *Eleutherodactylus coqui* 520–523.
- Revillagigedo Is., *Priacanthus (alalaua, boops)* 275–284.
- South Africa, *Leptobotula n. gen. breviventralis n. sp.* 166–170.
- South Carolina, *Sceloporus undulatus* 1–18; *Etmopterus (cyclosquamus n. sp., crossotus, microstomus, rimosus)* 140–156; *Rana sphenoccephala* 249–253; *Pseudemys scripta* 691–700; *Gambusia affinis* 937–945.
- Surinam, *Chelonia mydas* 798–802.
- Taiwan, *Bufo bankorensis* 561–579.
- Tennessee, *Sternotherus (minor, odoratus)* 285–294; *Etheostoma (lynceum, zonale)* 832–836.
- Texas, *Sceloporus undulatus* 1–18; *Etheostoma spectabile* 18–30; *Gambusia (affinis, nobilis)*, *Fundulus zebrinus*, *Cyprinodon variegatus*, *Percina macrolepidota* 86–91; *Hybopsis aestivalis*, *Nocomis leptoccephalus*, *Notemigonus crysoleucas*, *Notropis (amabilis, buchani, camurus, longirostris, lutrensis, oxyrhynchus, potteri, shumardi, venustus)*, *Pimephales notatus*, *Carpodius carpio*, *Ictalurus punctatus*, *Noturus gyrinus*, *Gambusia affinis*, *Lepomis (humilis, macrochirus)*, *Percina sciera*, *Mugil cephalus* 226–231; *Etheostoma whipplei* 631–641; *Ictalurus (lupus, punctatus)* 646–652; *Hemidactylus turcicus* 956–962; *Holbrookia propinqua* 980–986; *Thamnophis marcianus* 999–1001; *Sceloporus undulatus* 1012–1014.
- Thailand, *Oryzias mekongensis* 473–478.
- U.S.S.R., *Bufo g. gargaryans* 561–579.
- Utah, *Sceloporus undulatus* 1–18; *Rana pretiosa* 927–936.
- Venezuela, *Pygocentrus notatus*, *Pristobrycon striolatus*, *Serrasalmus rhombeus* 193–196; *Pimelodina flavipinnis* 653–672; *Oloolygon (danae n. sp., exigua n. sp.)* 864–870.
- Vermont, *Eurycea bislineata* 96–108; *Rana septentrionalis* 239–244.
- Virgin Is., *Priacanthus cruentatus* 275–284; *Dermochelys coriacea* 796–798; *Chelonia mydas* 798–802.
- Virginia, *Eurycea bislineata* 96–108; *Etmopterus microstomus* 140–156; *Chelydra serpentina*, *Malaclemys terrapin*, *Kinosternon subrubrum*, *Sternotherus odoratus*, *Clemmys guttata*, *Trionyx (muticus, spiniferus)*, *Chrysemys picta* 741–756; *Mustelus canis* 829–832; *Opsanus tau* 910–915.
- Washington, *Crotalus viridis oregonus* 816–820; *Rana (aurora, cascadae, pretiosa)* 927–936.
- West Virginia, *Notropis (s. spilopterus, s. hypsiosomatus)* 122–130; *Sternotherus odoratus* 285–294.
- Wisconsin, *Notropis (s. spilopterus, s. hypsiosomatus)* 122–130; *Rana septentrionalis* 239–244; *Necturus maculosus* 612–618; *Etheostoma zonale* 832–836.
- Wyoming, *Rana pretiosa* 927–936.
- Zimbabwe, *Crocodylus niloticus* 332–341.
- GROWTH, *Sceloporus (undulatus, clarki)* (seasonal rates) 1–18; *Etheostoma spectabile* (egg size relat. to growth rates of hatchlings) 18–30; *Bufo woodhousei*, *Hyla andersoni* (effect of low pH on larval growth) 454–466; *Gambusia affinis* (diff. in rate between adult males & females) 534–536; *Etheostoma juliae* (age-classes, male-female diff.) 536–540; *Chelydra serpentina* (salt marsh adap. for hatchling growth in saline water) 741–756; *Nerodia sipedon insularum* (ann. growth rate, sex. dimorphism) 757–772; *Chelonia mydas* (growth curve) 798–802; *Hemidactylus turcicus* (rates) 956–962; *Squatina californica* (rates in embryos) 987–994.



- HABITAT**, *Sceloporus (undulatus, clarki)* (hab. utilization) 1-18; (effects on resource partitioning in fishes) 352-388; *Rana utricularia*, *Limnaeodius ocularis*, *Pseudacris ornata*, *Bufo terrestris* (interspecific & size diff. in larval hab. use) 408-423; *Etheostoma (whipplei, radiosum)* 631-641; *Pseudemys scripta* (diff. hab. as dispersal barriers) 691-700; *Desmognathus (quadramaculatus, monticola, fuscus, ochrophaeus)* (affects community compos., exp. manip. in field) 731-741; *Cnemidophorus lemniscatus* (effect of microhab. on popn. dens.) 804-807; *Etheostoma (lynceum, zonale)* (spp. diff.) 832-836; *Phycis chesleri* (depth, temp.) 886-891; *Gambusia affinis* (hab. alter. affects gen. struct. of popns.) 937-945; *Acris crepitans* (color morph correlation with hab.) 995-999.
- HERMAPHRODITISM**, *Amphiprion clarkii* (sequential) 135-139; *Liopsetta glacialis* (1st report for sp., mixed gonad) 1024.
- HISTOLOGY**, *Iguana iguana* (skeletal muscle) 324-332; *Opsanus tau* (metab. chemicals in sonic muscle) 910-915.
- HOME RANGE**, *Amphiprion clarkii*, *Centropyge tibicen*, *Lactoria diaphana*, *Dendrochirus zebra*, *Plectrohynchus pictus* 135-139; *Ambystoma texanum* (absence of territor.) 725-730.
- HYBRIDIZATION**, *Hyla (andersonii × cinerea, andersonii × femoralis)* (natural hybrids) 70-76; *Eurycea (bislineata × n. sp.)* (of cryptic species) 96-108; *Morone (chrysops × americana)* (detect. by electrophoresis) 196-199; *Ictalurus (lupus, punctatus)* (electrophor. & morphomet. data) 646-652; *Tilapia (zilli, mariae)* (introduced popns.) 903-909; *Gila elegans* (*G. elegans × G. cypha*) 1021-1023.
- LACHRYMAL GLANDS**, *Dermochelys coriacea*, *Caretta caretta*, *Chelonia mydas*, *Eretmochelys imbricata* (compar. salt gland function) 247-249.
- LARVAE**, *Conger oceanicus* (capture of individual in tide pool) 222-223; *Rana utricularia*, *Limnaeodius ocularis*, *Pseudacris ornata*, *Bufo terrestris* (interspecific & size diff. in hab. use) 408-423; *Rana microdisca finchi* (transport on back of male) 523-525; *Hyla regilla* (feeding beh.) 802-804; *Hyla smithii* (described) 1004-1006; *Osteopilus septentrionalis* (cannibalism during metamorph.) 1007-1009.
- LATERAL LINE**, *Cottus bairdi* (response to prey) 91-96.
- LIFE HISTORY**, *Sceloporus (undulatus, clarki)* (including hab. effects in *S. undulatus*) 1-18; *Hexanchus griseus* 131-135; *Nerodia sipedon insularum* 757-772; *Saurida (undusquamis, longimanus, sp. 1, sp. 2)* (short generation times) 876-885; *Hemidactylus turcicus* (aspects relat. to successful colonizing) 956-962.
- LIGHT**, *Etheostoma spectabile* (photoper. during incubation affects fitness of hatchlings from var. egg sizes) 18-30.
- LOCOMOTION**, *Scomber japonicus* (effects of temp. on swimming speed) 39-44; *Gila bicolor mohavensis* (met. rates at var. temps. and swimming speeds) 45-52; *Caretta caretta* (swimming & diving techniques in hatchlings) 53-57; *Rhineura floridana*, *Elaphe obsoleta*, *Nerodia fasciata*, *Acrochordus javanicus*, *Cerberus rhyhops*, *Crotalus cerastes* (kinematics, terrest. & aquat.) 915-927; *Thamnophis marcianus* (affected by food intake) 999-1001.
- METHODS**, *Etheostoma proeliare* (marking with fluorescent plastic chips for small fishes) 544-546; (polarized light aids fin ray counts) 825-826; *Gopherus agassizii* (meth. to improve capture of juvs.) 1010-1012.
- MIGRATION**, *Ambystoma macrodactylum columbianum* (breeding migration) 398-408; *Atelopus varius* (homing after exp. displacement) 438-444; *Gambusia affinis* (diff. dispersal patterns of juvs. & adults) 937-945; *Salvelinus alpinus* (spawning & feeding) 945-955.
- MORPHOLOGY**, *Anolis (distichus dominicensis, d. floridanus)* subspecific intergrad. in scale charc.) 76-86; *Heterandria (formosa, bimaculata, jonesi)*, *Gambusia (affinis, nobilis, yucatanana)*, *Poeciliopsis sp.*, *Xiphophorus (maculatus, helleri)*, *Priapella compressa*, *Poecilia (butleri, reticulata, latipinna)*, *Belonesox belizanus*, *Profundulus (labialis, punctatus)*, *Fundulus zebrinus*, *Cyprinodon variegatus*, *Percina macrolepida* (of scales) 86-91; *Notropis (s. spilopterus, s. hypsismatus)* (truss anal. of shape) 122-130; *Evermannichthys (metzelaari, spongicola)* (merist. & morph. var.) 157-165; *Notropis (calientis, amecae n. sp., aulidion n. sp.)* (key to *Notropis* s. of 24°N) 170-183; *Pygocentrus notatus*, *Pristigobryon striolatus*, *Serrasalminus rhombus* (relat. to feeding) 193-196; *Hyposblennius (ionthas, hentsi)*, *Hypleurochilus geminatus*, *Parablennius marmoratus* (teeth and jaws) 207-213; *Rhinichthys cataractae* (diff. between popns. isol. only 40 yrs) 219-221; *Manacopus fal-cifer* (homol. of pelvic fins and priapium) 305-310; *Embiotoca (jacksoni, lateralis)* (pharyngeal jaw apparatus, bones & muscles) 311-323; (jaw muscle in higher teleosts) 705-713; *Rana (a. aurora, a. draytonii, boylii, cascadeae, muscosa, pretiosa)* (anat. of vocal sacs) 927-936; *Cichlasoma citrinellum* (lower jaw, head shape, relat. to feeding) 1028-1030.
- NESTING**, *Pituophis melanoleucus* (nest site selection) 116-121; *Sceloporus aeneus* 232-233; *Caretta caretta* (power plant effects) 813-816.
- NOMENCLATURE**, (keys for identifying fishes described by Cuvier and Valenciennes) 216-219.
- PALEONTOLOGY**, *Sauromalus sp.*, *Gambelia wislizeni*, *Sceloporus sp.*, *Phrynosoma sp.*, *Eumeces gil-*

*bertii*, *Cnemidophorus*?, *Coleonyx* sp. (late Wisconsin lizard fauna) 244–246; *Heterodon* (*nasicus*, *platyrhinos*), *Coluber* or *Masticophis*, *Elaphe vulpina*, *Lampropeltis* (*getulus*, *triangulum*), *Pituophis melanoleucus*, *Thamnophis* cf. *radix*, *Nerodia* sp. indet. (early Illinoian, Okla.) 811–812.

**PHEROMONES**, *Ictalurus nebulosus* (aggression suppressing) 540–544.

**PHYSIOLOGY**, *Gila bicolor mohavensis* (met. rates, temp. and salin. toler.) 45–52; *Dermochelys coriacea*, *Caretta caretta*, *Chelonia mydas*, *Eretmochelys imbricata* (salt excretion) 247–249; *Iguana iguana* (skeletal muscle) 324–332; *Ctenotus taeniolatus* (season. pattern of lipid storage and metab.) 445–453; *Sauromalus hispidus* (high extracell. K<sup>+</sup> affects body temp. & feeding) 518–520; *Carcharodon carcharias*, *Isurus oxyrinchus*, *Carcharhinus plumbeus*, *obscurus*, *Galeocerdo cuvieri*, *Prionace glauca*, *Sphyrna lewini*, *Alopius vulpinus* (hematol. of endotherms vs ectotherms) 700–705; *Gopherus polyphemus* (O<sub>2</sub> & CO<sub>2</sub> in burrows) 787–795; *Dermochelys coriacea* (met. rate, food requirements) 796–798; *Opsanus tau* (fast sonic muscle) 910–915.

**POPULATIONS**, *Sceloporus undulatus, clarki* (compar. demographics) 1–18; *Cyclura nubila* (popn. energy expenditure est. from body temp. data) 65–69; *Anolis* (*distichus dominicensis*, *d. floridanus*) (subspecific intergrad. following colonization) 76–86; *Echidna zebra*, *Aulostomus chinensis*, *Fistularia commersonii*, *Mugil cephalus, curema*, *Priacanthus* (*alalaua*, *arenatus*, *boops*, *cruentatus*), *Coryphaena hippurus*, *Gnathanodon speciosus*, *Zanclus cornutus*, *Alutera scripta*, *Arothron* (*hispidus*, *meleagris*), *Diodon holocanthus* (electrophor. compar. between trans-Pacific popns.) 275–284; *Ambystoma macrodactylum columbianum* (demography & migration) 398–408; *Acrochordus arafurae* (density of a low-energy specialist) 424–437; *Pseudemys scripta* (electroph. diff. between popns. sep. by diff. hab.) 691–700; *Nerodia sipedon insularum* (size, struct., diff. between island & mainland) 757–772; *Cnemidophorus lemniscatus* (ecol. factors affecting popn. density) 804–807. *Saurida* (*undosquamis*, *longimanus*, sp. 1, sp. 2) (size & spp. compos. changes with fishing) 876–885; *Gambusia affinis* (gen. struct.) 937–945.

**PREDATION**, *Aprasia* (*parapulchella*, *pulchella*, *repens*, *striolata*), *Delma* (*fraseri*, *inornata*, *nasuta*), *Lialis* (*burtonis*, *ficari*), *Pygopus* (*lepidopodus*, *nigriceps*) (compar. prey types) 30–39; *Cottus bairdi* (lat. line receptivity to prey stim.) 91–96; *Ammodytes hexapterus* (intensified crepuscular pred. pressure) 223–226; *Acrochordus arafurae* (selective effects of human pred.) 238–239; *Acrochordus arafurae* (piscivory in a low-energy specialist) 424–437; *Darlingtonia haetiana* (single prey genus) 529–531; *Nerodia sipedon insularum* (evid. of low pred. level) 757–772; *Crotalus viridis oreganus* (strike effectiveness) 816–820; *Phrynosoma* (*cornutum*, *modestum*) (death rate by pred.) 820–824; *Clemmys in-*

*sculpta* (stomping for earthworms) 1001–1004; *Hyla smithii* (effect of tail color) 1004–1006; *Osteopilus septentrionalis* (larval cannibalism during metamorph.) 1007–1009.

**REPRODUCTION**, *Sceloporus undulatus, clarki* (reprod. season & clutch freq., mass) 1–18; *Aprasia* (*parapulchella*, *pulchella*, *repens*, *striolata*), *Delma* (*fraseri*, *inornata*, *nasuta*), *Lialis* (*burtonis*, *ficari*), *Pygopus* (*lepidopodus*, *nigriceps*) (seasonal., clutch size) 30–39; *Hyla* (*andersonii* × *cinerea*, *andersonii* × *femorialis*) (infertility of hybrids) 70–76; *Padogobius martensi* (sounds made during courtship and terr. defense) 213–216; *Sceloporus aeneus* (nesting and oviposition) 232–233; *Rana septentrionalis* (requirements of reprod. limit s. distribution) 239–244; *Rana sphenoccephala* (low temp. correlated with communal egg-laying) 249–253; *Poecilia latipinna* (interbrood interval) 295–304; *Fundulus heteroclitus* (season. patterns var. with size) 342–351; *Ambystoma macrodactylum columbianum* (breeding migration) 398–408; *Acrochordus arafurae* (in a low-energy specialist) 424–437; *Ctenotus taeniolatus* (relat. to season. pattern of lipid storage) 445–453; *Poecilia latipinna* (fertility advertisement by females) 467–472; *Eleuthero-dactylus coqui* (season. effect of temp.) 520–523; *Etheostoma juliae* (seasonal. of gonads & second. charc.) 536–540; *Hyla crucifer* (satellite males) 719–724; *Nerodia sipedon insularum* (season., size-dependence, fecundity) 757–772; *Lygodactylus klugei*, *Phyllorhynchus pollicaris*, *Gymnodactylus geckoides*, *Hemidactylus mabouia* (size at sex. mat., invariable clutch size, season. cycle) 773–786; *Aghistrodon contortrix* (sperm storage, mult. paternity in lab) 807–811; *Caretta caretta* (power plant affects nesting) 813–816; *Saurida* (*undosquamis*, *longimanus*, sp. 1, sp. 2) (age & size at sex. mat.) 876–885; *Salvelinus alpinus* (spawning interval relat. to energy content) 945–955; *Hemidactylus turcicus* (aspects relat. to successful colonizing) 956–962; *Sceloporus jarrovi* (lower body temp. of preg. females) 971–979; *Holbrookia propinqua* (female breeding colors) 980–986; *Squatina californica* (ann. cycle, sex. matur., fecund.) 987–994; *Aphredoderus sayanus* (branchial brooding) 1030–1031.

**RESPIRATION**, *Gopherus polyphemus* (O<sub>2</sub> & CO<sub>2</sub> in burrows) 787–795; *Dermochelys coriacea* (resp. rate of hatchlings) 796–798.

**SALINITY**, *Gila bicolor mohavensis* (tolerance levels) 45–52; *Chelydra serpentina*, *Malaclemys terrapin*, *Kinosternon subrubrum*, *Sternotherus odoratus*, *Clemmys guttata*, *Trionyx* (*muticus*, *spiniferus*), *Chrysemys picta* (adapt. to estuarine water) 741–756.

**SEX**, *Lialis* (*burtonis*, *ficari*) (skewed sex ratios) 30–39; *Phrynosoma* (*modestum*) (ratio) 886–891.

**SEXUAL DIMORPHISM**, *Sceloporus undulatus, clarki* (in size) 1–18; *Aprasia* (*parapulchella*, *pulchella*,


- repens*, *striolata*), *Delma* (*fraseri*, *inornata*, *nasuta*), *Lialis* (*burtonis*, *jacari*), *Pygopus* (*lepidopodus*, *nigriceps*) (in size) 30–39.
- SIZE, *Sceloporus* (*undulatus*, *clarki*) at matur., sex. dimorph.) 1–18; *Etheostoma spectabile* (egg size: seasonal var., relat. to hatchling size and fitness) 18–30; *Aprasia* (*parapulchella*, *pulchella*, *repens*, *striolata*), *Delma* (*fraseri*, *inornata*, *nasuta*), *Lialis* (*burtonis*, *jacari*), *Pygopus* (*lepidopodus*, *nigriceps*) (sex. dimorph.) 30–39; *Fundulus heteroclitus* (var. in seasonal reprod. patterns with size) 342–351; *Hyla crucifer* (calling compar. to non-calling males, relative amplexus success) 719–724; *Rana utricularia*, *Limnaeodius ocularis*, *Bufo terrestris* (intraspecific size diff. of larvae affects hab. use) 408–423; *Nerodia sipidon insularum* (season. diff., sexual dimorph., size at matur., relat. to weight) 757–772; *Lygodactylus klugei*, *Phyllopleopus pollicaris*, *Gymnodactylus gekooides*, *Hemidactylus mabouia* (sex. dimorph., invariable clutch size affects female size) 773–786; *Salvelinus alpinus* (of reprod. individuals) 945–955; *Hemidactylus turcicus* (season. changes in size classes) 956–962; *Squatina californica* (at sex. matur.) 987–994.
- SURVIVAL, *Sceloporus* (*undulatus*, *clarki*) (egg to yearling, adults between sexes over 3 years) 1–18; *Hemidactylus turcicus* (of diff. age classes) 956–962.
- SYSTEMATICS, *Heterandria* (*formosa*, *bimaculata*, *jonesi*), *Gambusia* (*affinis*, *nobilis*, *yucatanana*), *Poeciliopsis* sp., *Xiphophorus* (*maculatus*, *helleri*), *Priapella compressa*, *Poecilia* (*butleri*, *reticulata*, *latipinna*), *Belonesox belizanus*, *Profundulus* (*labialis*, *punctatus*), *Fundulus zebrinus*, *Cyprinodon variegatus*, *Percina macrolepidia* (lepidonts on scales as taxon. char.) 86–91; *Eurycea* (n. sp., b. *bilineata*, b. *wilderae*, b. *cirrigera*, b. *rivicola*) (electroph. detection of cryptic species) 96–108; *Notropis* (s. *spiloteris*, s. *hyposomatus*) (invalid subsp.) 122–130; *Etopus* (*cycloquamus* n. sp., *crossotus*, *microstomus*, *rimosus*) 140–156; *Evermannichthys* (*metzelaari*, *spongicola*) (status of unique specimens) 157–165; *Leptobrotula* n. gen. *breviventralis* n. sp. 166–170; *Notropis* (*calientis*, *amecae* n. sp., *aulidion* n. sp.) 170–183; *Cyprinodon pachecephalus* n. sp. 184–192; *Kinosternon* (*bauri*, *subrubrum*, *herrerai*, *hirtipes*, *sonoriense*, *flavescens*, *alamosae*, *integrum*, *scorpioides*, *oaxacae*, *acutum*, *leucostomum*), *Staurotypus triporcatus*, *Claudius angustatus*, *Sternotherus* (*odoratus*, *carinatus*, *depressus*, *minor*) (electroph. compar. of family) 285–294; *Oryzias melanocephalus* (Mekong basin, Thailand) 473–478; *Ceratias* (*tentaculatus*, *holboellii*, *uranoscopus*), *Cryptopsaras couesi* (rev. of family) 479–493; *Stomias* (*brevibarbatulus*, *danae*, *longibarbatulus*, *pacificus*, *colubrinus*, *lampropeltis*, *affinis*, *nebulosus*, *atriventer*, *boa*, *gracilis*) (phylogen. anal., including evid. for synonymy of *Macrostomias*) 494–503; *Eleutherodactylus* (*andi*, *fitzingeri*, *longirostris*, *raniformis*, *cuaquero*, *melanostictus*, *monnichorum*, *rayo*, *crassidigitus*, *italamancae*) (cladogram [PAUP program] & phylogen. class. based on allozymes, karyol., morphol.) 503–511; *Bufo* (*tibetanus*, *minshanicus*, *j. japonicus*, *j. formosus*, *torrenticola*, *g. gargazianus*, *g. miyakonis*, *g. popei*, *bankorensis*, *andrewsi*) (geogr. var., 3 spp. synonymized) 561–579; *Centropomus* (*pectinatus*, *medius*, *undecimalis*, *nigrescens*, *viridis*, *poei*, *parallelus*, *mexicanus*, *unionensis*, *armatus*, *robalius*, *ensiferus*) (rev. of genus, incl. key) 579–611; *Osornophryne* (*talipes* n. sp., *percrassa*, *bufoniformis*) 618–622; *Amolops* (*amoropalampus* n. sp., *poecilus*, *whiteheadi*, *orphnocnemis* n. sp., *phaeomerus*, *jerboa*, *macrophthalmus* n. sp.) (new spp., key to group) 623–630; *Etheostoma* (*whipplei*, *radiosum*) 631–641; *Apogon chalcus* n. sp. 641–645; *Ictalurus* (*lupus*, *punctatus*) (electroph. & morphomet. data) 646–652; *Aguarunichthys torosus*, *Pimelodina flavipinnis*, *Calophysus macropterus*, *Lucioperca latipinna*, *Pini-rampus pini-rampu* (new genus, rev. of *Pimelodina*) 653–672; *Pterois* (*volitans*, *miles*) (valid. of sp.) 686–690; *Clupea* (*harengus*, *pallasii*) (elev. to sp. status) 714–719; *Etheostoma* (*lynceum*, *zonale*) (elev. to sp. status) 832–836; *Simuliparna* n. gen. *hermani* (Cape Verde Archipelago) 857–863; *Olophryne* (*danae* n. sp., *exigua* n. sp.) (Venezuela) 864–870; *Pseudotropheus barlowi* n. sp. (Lake Malawi) 870–875; *Anchoa* (*starksi*, *marinii*, *naso*, *filifera*, *lyolepis*, *nasus*, *nasuta*, *longipinna*), *Anchoviella* n. subgen., *Anchoviella elongata* (spp. groups of *Anchoa*) 891–902; *Eristocophis* (*mcnamahoni*, *arietans*, *nasicornis*, *gabonica*), *Causus maculatus*, *Echis carinatus*, *Micrurus nigrocinctus* (electrophor. anal., PAUP program).
- TEMPERATURE, *Etheostoma spectabile* (incubation effects on fitness of hatchlings from var. egg sizes) 18–30; *Scomber japonicus* (effects on swimming speed, lethal limits) 39–44; *Gila bicolor mohavensis* (toler. levels, met. rates at var. temps.) 45–52; *Ctenopus* (*atlas*, *regius*, *robustus*, *taeniolatus*, *uber*), *Egernia* (*striolata*, *whitii*), *Eremiascincus* (*fasciolatus*, *richardsoni*), *Hemiergis* (*decreciensis*, *peroni*), *Leiolopisma entrecasteauxii*, *Sphenomorphus* (*kosciuskoii*, *quoyi*, *tympanum*), *Tiliqua* (*cassurinae*, *rugosa*) (tolerance limits, preferred body temp.) 57–64; *Cyclura nubila* (thermoreg. in free-ranging) 65–69; *Hyla crucifer* (affects color change) 109–115; *Etopus* (*cycloquamus* n. sp., *crossotus*, *microstomus*, *rimosus*) (eff. on distr.) 140–156; *Cyprinodon pachecephalus* n. sp. (hab. 44°C thermal springs) 184–192; *Sceloporus aeneus* (of natural nests) 232–233; *Rana sphenoccephala* (low temp. correlated with communal egg-laying) 249–253; *Poecilia latipinna* (affects interbrood interval) 295–304; *Sauromalus hispidus* (high extracell. K<sup>+</sup> affects thermoreg.) 518–520; *Eleutherodactylus coqui* (effect on rate of develop. in terrestrial-breeding frog) 520–523; *Necturus maculosus* (diel & season. var. in beh. thermoreg.) 612–618; *Carcharodon carcharias*, *Isurus oxyrinchus*, *Carcharhinus* (*plumbeus*, *obscurus*), *Galeocerdo cuvieri*, *Prionace glauca*, *Sphyrna lewini*, *Alopias vulpinus* (hematol. of endotherms

vs ectotherms) 700-705; *Gopherus polyphemus* (in burrows) 787-795; *Phycis chesteri* (preference) 886-891; *Sceloporus jarrovi* (lower body temp. of preg. females) 971-979; *Gopherus agassizii* (toler. of juvs. to lower air temp.) 1010-1012; *Sceloporus undulatus* (eff. of cold on eggs) 1012-1014.

TRUSS ANALYSIS, *Notropis* (*s. spilopterus*, *s. hypsisomatus*) (detects invalidity of subspp.) 122-130; *Tilapia* (*zilli*, *mariae*) (ident. of hybrids) 903-909.

URINARY BLADDER, (presence or absence in lizards) 512-517.

VARIATION, *Notropis* (*s. spilopterus*, *s. hypsisomatus*) (geogr. var. in shape & merist.) 122-130; *Evermannichthys* (*metzelaari*, *spongicola*) (geogr.) 157-165; *Bufo* (*tibetanus*, *minshanicus*, *j. japonicus*, *j. formosus*, *torrenticola*, *g. gargarizans*, *g. miyakonis*, *g. popei*, *bankorensis*, *andrewsi*) (geogr.) 561-579; *Etheostoma* (*whipplei*, *radiosum*) (geogr.) 631-641.





# CONTENTS—1986, Nos. 1-4

ADKINS, M. DALE (see Seidel, Michael E.)	
ALFORD, ROSS A. Habitat use and positional behavior of anuran larvae in a northern Florida temporary pond	408
AMEMIYA, CHRIS T. AND JOHN R. GOLD. Chromomycin A <sub>3</sub> stains nucleolus organizer regions of fish chromosomes	226
AMEMIYA, CHRIS T., STEVEN W. KELSCH, FRED S. HENDRICKS AND JOHN R. GOLD. The karyotype of the Mexican blindcat, <i>Prietella phreatophila</i> Cartanza (Ictaluridae)	1024
ANDERSON, KAREN AND PAUL E. MOLER. Natural hybrids of the pine barrens treefrog, <i>Hyla andersonii</i> with <i>H. cinerea</i> and <i>H. femoralis</i> (Anura, Hylidae): morphological and chromosomal evidence	70
ANDERSON, JOHN F. (see Ultsch, Gordon R.)	
BELS, VINCENT L. Analysis of the display-action-pattern of <i>Anolis chlorocyanus</i> (Sauria: Iguanidae)	963
BENESKI, JOHN T., JR., EDWARD J. ZALISKO AND JOHN H. LARSEN, JR. Demography and migratory patterns of the eastern long-toed salamander, <i>Ambystoma macrodactylum columbianum</i>	398
BENNETT, ALBERT F. AND HENRY JOHN-ALDER. Thermal relations of some Australian skinks (Sauria: Scincidae)	57
BERRY, KRISTIN H. AND FREDERICK B. TURNER. Spring activities and habits of juvenile desert tortoises, <i>Gopherus agassizii</i> , in California	1010
BEUCHAT, CAROL A. Phylogenetic distribution of the urinary bladder in lizards	512
BEUCHAT, CAROL A. Reproductive influences on the thermoregulatory behavior of a live-bearing lizard	971
BJORNDALE, KAREN A. Effect of solitary vs group feeding on intake in <i>Pseudemys nelsoni</i>	234
BLEM, CHARLES R. (see Fine, Michael L.)	
BOLTZ, J. M. AND J. R. STAUFFER, JR. Branchial brooding in the pirate perch, <i>Aphredoderus sayanus</i> (Gilliams)	1030
BRASWELL, ALVIN L. (see Wiley, John E.)	
BURDICK, BOB D. (see Kaeding, Lynn R.)	
BURGER, JOANNA AND ROBERT T. ZAPPALORTI. Nest site selection by pine snakes, <i>Pituophis melanoleucus</i> , in the New Jersey Pine Barrens	116
BURGESS, GEORGE H. (see Gilbert, Carter R.)	
CAILLIET, GREGOR M. (see Natanson, Lisa J.)	
CALDWELL, JANALEE P. Selection of egg deposition sites: a seasonal shift in the southern leopard frog, <i>Rana sphenoccephala</i>	249
CALDWELL, JANALEE P. A description of the tadpole of <i>Hyla smithii</i> with comments on tail coloration	1004
CANNATELLA, DAVID C. A new species of <i>Osornophryne</i> (Anura, Bufonidae) from the Andes of Ecuador	618
CARBONE, P. (see Vitturi, R.)	
CARR, JOHN E. (see Carr, Marilyn G.)	
CARR, MARILYNN G. AND JOHN E. CARR. Characterization of an aggression suppressing pheromone in the juvenile brown bullhead ( <i>Ictalurus nebulosus</i> )	540
CATALANO, E. (see Vitturi, R.)	
CAVENDER, TED M. (see Schaefer, Scott A.)	
CHERNOFF, BARRY AND ROBERT RUSH MILLER. Fishes of the <i>Notropis calientis</i> complex with a key to the southern shiners of Mexico	170
CHESSER, RONALD K. (see Kennedy, Phyllis K.)	
CHRISTIAN, KEITH A., ILEANA E. CLAVIJO, NANCY CORDERO-LOPEZ, EVELYN E. ELIAS-MALDONADO, MIGUEL A. FRANCO, MABEL V. LUGO-RAMIREZ AND MARISOL MARENGO. Thermoregulation and energetics of a population of Cuban iguanas ( <i>Cyclura nubila</i> ) on Isla Maguayes, Puerto Rico	65
CHRISTIAN, KEITH A., C. RICHARD TRACY AND WARREN P. PORTER. The effect of cold exposure during incubation of <i>Sceloporus undulatus</i> eggs	1012
CLARK, JENNIFER B. (see Helfman, Gene S.)	
CLAVIJO, ILEANA E. (see Christian, Keith A.)	
CLOUGH, WENDI (see Davenport, John)	
COOPER, WILLIAM E., JR. Chromatic components of female secondary sexual coloration: influence on social behavior of male keeled earless lizards ( <i>Holbrookia propinqua</i> )	980
CORDERO-LOPEZ, NANCY (see Christian, Keith A.)	
COURTENAY, WALTER R., JR. (see Taylor, Jeffrey N.)	
CRUMP, MARTHA L. Homing and site fidelity in a neotropical frog, <i>Atelopus varius</i> (Bufonidae)	438



CRUMP, MARTHA L. Cannibalism by younger tadpoles: another hazard of metamorphosis .....	1007
DAVENPORT, JOHN AND WENDI CLOUGH. Swimming and diving in young loggerhead sea turtles ( <i>Caretta caretta</i> L.) .....	53
DILLAMAN, RICHARD M. (see Lindquist, David G.)	
DRUMMOND, KATHLEEN A. (see Fine, Michael L.)	
DUELLMAN, WILLIAM E. Two new species of <i>Ololygon</i> (Anura: Hylidae) from the Venezuelan Guyana .....	864
DUNHAM, ARTHUR E. (see Tinkle, Donald W.)	
DUNSON, WILLIAM A. (see Freda, Joseph)	
DUNSON, WILLIAM A. Estuarine populations of the snapping turtle ( <i>Chelydra</i> ) as a model for the evolution of marine adaptations in reptiles .....	741
DUTIL, J. D. Energetic constraints and spawning interval in the anadromous Arctic charr ( <i>Salvelinus alpinus</i> ) .....	945
EBERT, DAVID A. Biological aspects of the sixgill shark, <i>Hexanchus griseus</i> .....	131
ELIAS-MALDONADO, EVELYN E. (see Christian, Keith A.)	
EMERY, SCOTT H. Hematological comparisons of endothermic vs ectothermic elasmobranch fishes .....	700
ERNST, ROBERT G. (see Proffit, C. Edward)	
ETNIER, DAVID A. AND WAYNE C. STARNES. <i>Etheostoma lynceum</i> removed from the synonymy of <i>E. zonale</i> (Pisces, Percidae) .....	832
EVANS, J. E. (see Scribner, K. T.)	
FARR, JAMES A. AND JOSEPH TRAVIS. Fertility advertisement by female sailfin mollies, <i>Poecilia latipinna</i> (Pisces: Poeciliidae) .....	467
FELDMETH, C. ROBERT (see McClanahan, Lon L.)	
FINE, MICHAEL L., KEITH R. PENNYPACKER, KATHLEEN A. DRUMMOND AND CHARLES R. BLEM. Concentration and location of metabolic substrates in fast toadfish sonic muscle .....	910
FINK, SARA V. (see Fink, William L.)	
FINK, WILLIAM L. AND SARA V. FINK. A phylogenetic analysis of the genus <i>Stomias</i> , including the synonymization of <i>Macrostomias</i> .....	494
FORD, NEIL B. AND GLADIS A. SHUTTLESWORTH. Effects of variation in food intake on locomotory performance of juvenile garter snakes .....	999
FORESTER, DON C. AND DAVID V. LYKENS. Significance of satellite males in a population of spring peepers ( <i>Hyla crucifer</i> ) .....	719
FRANCO, MIGUEL A. (see Christian, Keith A.)	
FRANKE, CARLOS ROBERTO (see Magnusson, William E.)	
FRASER, THOMAS H. AND JOHN E. RANDALL. A new species of the cardinalfish genus <i>Apogon</i> from Easter Island .....	641
FRAZER, NAT B. AND ROSAMUND C. LADNER. A growth curve for green sea turtles, <i>Chelonia mydas</i> , in the U.S. Virgin Islands, 1913-14 .....	798
FREDA, JOSEPH AND WILLIAM A. DUNSON. Effects of low pH and other chemical variables on the local distribution of amphibians .....	454
GARCIA, CARMEN (see Machado-Allison, Antonio)	
GIBBONS, J. W. (see Scribner, K. T.)	
GILBERT, CARTER R. AND GEORGE H. BURGESS. Variation in western Atlantic gobiid fishes of the genus <i>Evermannichthys</i> .....	157
GILLINGHAM, JAMES C. (see Schuett, Gordon W.)	
GLEESON, TODD T. AND JON M. HARRISON. Reptilian skeletal muscle: fiber-type composition and enzymatic profile in the lizard, <i>Iguana iguana</i> .....	324
GOLD, JOHN R. (see Amemiya, Chris T.)	
GOLD, JOHN R. (see Amemiya, Chris T., et al.)	
GOLDBERG, STEPHEN R., WILLIAM T. YASUTAKE AND ROBIN L. WEST. A hermaphroditic Arctic flounder, <i>Liopsetta glacialis</i> (Pleuronectidae), from Alaska .....	1024
GONGORA, GUILLERMO LARA (see Guillelte, Louis J., Jr.)	
GORMAN, WENDY L. Pattern of color polymorphism in the cricket frog, <i>Acris crepitans</i> , in Kansas .....	995
GOSLINE, WILLIAM A. Jaw muscle configuration in some higher teleostean fishes .....	705
GOTTFRIED, MICHAEL D. Developmental transition in feeding morphology of the midas cichlid .....	1028
GRANT, W. STEWART. Biochemical genetic divergence between Atlantic, <i>Clupea harengus</i> , and Pacific, <i>C. pallasi</i> , herring .....	714
GRAUNKE, BARBARA J. (see Proffit, C. Edward)	
GUILLETTE, LOUIS J., JR. AND GUILLERMO LARA GONGORA. Notes on oviposition and nesting in the high elevation lizard, <i>Sceloporus aeneus</i> .....	232
GUNN, J. S. (see Thresher, R. E.)	
GUTTMAN, SHELDON I. AND ALVAN A. KARLIN. Hybridization of cryptic species of two-lined salamanders ( <i>Eurycea bislineata</i> complex) .....	96



HARDER, WILHELM. Polarized light as an aid in counting fin rays .....	825
HARRISON, JON M. (see Gleeson, Todd T.)	
HAYES, MARC P. (see Miyamoto, Michael M.)	
HAYES, MARC P. AND DANA M. KREMPELS. Vocal sac variation among frogs of the genus <i>Rana</i> from western North America .....	927
HEDEEN, STANLEY E. The southern geographic limit of the mink frog, <i>Rana septentrionalis</i> .....	239
HELFMAN, GENE S. AND JENNIFER B. CLARK. Rotational feeding: overcoming gape-limited foraging in anguillid eels .....	679
HENDERSON, ROBERT W. AND ALBERT SCHWARTZ. The diet of the Hispaniolan colubrid snake, <i>Darlingtonia haetiana</i> .....	529
HENDRICKS, FRED S. (see Kelsch, Steven W.)	
HENDRICKS, FRED S. (see Amemiya, Chris T.)	
HENSLEY, DANNIE A. A new damselfish genus from the Cape Verde Archipelago based on <i>Glyphidodon (Parma) hermani</i> Steindachner, 1887 (Pisces: Pomacentridae) .....	857
HOBSON, EDMUND S. Predation on the Pacific sand lance, <i>Ammodytes hexapterus</i> (Pisces: Ammodytidae), during the transition between day and night in southeastern Alaska .....	223
HOEKSTRA, DENISE AND JOHN JANSSEN. Lateral line receptivity in the mottled sculpin ( <i>Cottus bairdi</i> ) ..	91
HOLLANDER, ROBERT R. Microanalysis of scales of poeciliid fishes .....	86
HOLMAN, J. ALAN. Snakes of the Berends local fauna (Pleistocene: Early Illinoian) of Oklahoma .....	811
HUDSON, DAVID M. AND PETER L. LUTZ. Salt gland function in the leatherback sea turtle, <i>Dermochelys coriacea</i> .....	247
HUGHES, AUSTIN L. Growth of adult mosquitofish <i>Gambusia affinis</i> in the laboratory .....	534
HUTCHISON, VICTOR H. AND KARLA K. SPIESTERSBACH. Diel and seasonal cycles of activity and behavioral thermoregulation in the salamander <i>Necturus maculosus</i> .....	612
HUTTON, J. M. Age determination of living Nile crocodiles from the cortical stratification of bone ...	332
INGER, ROBERT F., HAROLD K. VORIS AND PAUL WALKER. Larval transport in a Bornean ranid frog ...	523
IVERSON, JOHN B. (see Seidel, Michael E.)	
JACOBSON, L. D., R. L. TORBLAA AND R. H. MORMAN. Temporal variation in allelic frequencies of sea lamprey ammocoetes .....	199
JAEGER, ROBERT G. (see Martin, David L.)	
JAMES, PAUL W. AND CHARLES A. TABER. Reproductive biology and age and growth of the yoke darter, <i>Etheostoma juliae</i> .....	536
JANSSEN, JOHN (see Hoekstra, Denise)	
JAYNE, BRUCE C. Kinematics of terrestrial snake locomotion .....	915
JOHN-ALDER, HENRY (see Bennett, Albert F.)	
JONES, JEFF (see McClanahan, Lon L.)	
KAEDING, LYNN R., BOB D. BURDICK, PATRICIA A. SCHRADER AND WILLIAM R. NOONAN. Recent capture of a bonytail ( <i>Gila elegans</i> ) and observations on this nearly extinct cyprinid from the Colorado River	1021
KARDONG, KENNETH V. The predatory strike of the rattlesnake: when things go amiss .....	816
KARLIN, ALVAN A. (see Guttman, Sheldon I.)	
KASPER, LUIZ ALBERTO (see Magnusson, William E.)	
KATS, LEE B. AND RANDALL G. VAN DRAGT. Background color-matching in the spring peeper, <i>Hyla crucifer</i> .....	109
KAUFMANN, JOHN H. Stomping for earthworms by wood turtles, <i>Clemmys insculpta</i> : a newly discovered foraging technique .....	1001
KELLER, E. C., JR. (see Whitman, Pamela A.)	
KELSCH, STEVEN W. (see Amemiya, Chris T.)	
KELSCH, STEVEN W. AND FRED S. HENDRICKS. An electrophoretic and multivariate morphometric comparison of the American catfishes <i>Ictalurus lupus</i> and <i>I. punctatus</i> .....	646
KENNEDY, MICHAEL L. (see Kennedy, Phyllis K.)	
KENNEDY, PHYLLIS K., MICHAEL L. KENNEDY, EARL G. ZIMMERMAN, RONALD K. CHESSER AND MICHAEL H. SMITH. Biochemical genetics of mosquitofish. V. Perturbation effects on genetic organization of populations .....	937
KING, RICHARD B. Population ecology of the Lake Erie water snake, <i>Nerodia sipedon insularum</i> .....	757
KNEIB, RONALD T. Size-specific patterns in the reproductive cycle of the killifish, <i>Fundulus heteroclitus</i> (Pisces: Fundulidae) from Sapelo Island, Georgia .....	342
KNIGHT, LUTHER A., JR. (see Thompson, Kenneth W.)	
KREMPELS, DANA M. (see Hayes, Marc P.)	
LABAT, CHRISTOPHER P. (see Martin, David L.)	
LADNER, ROSAMUND (see Frazer, Nat B.)	
LARGE, HELEN L. (see Snelson, Franklin F., Jr.)	
LARSEN, JOHN H., JR. (see Beneski, John T., Jr.)	

LeCroy, Sara E. (see Proffit, C. Edward)	
Leslie, Andrew J., Jr. and Donald J. Stewart. Systematics and distributional ecology of <i>Etropus</i> (Pisces: Bothidae) on the Atlantic Coast of the United States with description of a new species	140
Liem, Karel F. The pharyngeal jaw apparatus of the Embiotocidae (Teleostei): a functional and evolutionary perspective	311
Lillywhite, Harvey (see Smits, Allan W.)	
Lindquist, David G. and Richard M. Dillaman. Trophic morphology of four western Atlantic blennies (Pisces: Blenniidae)	207
Lugo-Ramirez, Mabel V. (see Christian, Keith A.)	
Lutcavage, Molly and Peter L. Lutz. Metabolic rate and food energy requirements of the leatherback sea turtle, <i>Dermochelys coriacea</i>	796
Lutz, Peter L. (see Hudson, David M.)	
Lutz, Peter L. (see Lutcavage, Molly)	
Lykens, David V. (see Forester, Don C.)	
Macalusco, M. (see Vitturi, R.)	
Machado-Allison, Antonio and Carmen Garcia. Food habits and morphological changes during ontogeny in three serrasalmin fish species of the Venezuelan floodplains	193
Magnusson, William E., Carlos Roberto Franke and Luiz Alberto Kasper. Factors affecting densities of <i>Cnemidophorus lemniscatus</i>	804
Magtoon, Wichian (see Uwa, Hiroshi)	
Marengo, Marisol (see Christian, Keith A.)	
Marsh, Edie. Effects of egg size on offspring fitness and maternal fecundity in the orangethroat darter, <i>Etheostoma spectabile</i> (Pisces: Percidae)	18
Marshall, Joseph A. (see Whitman, Pamela A.)	
Martin, David L., Robert G. Jaeger and Christopher P. Labat. Territoriality in an <i>Ambystoma</i> salamander? Support for the null hypothesis	725
Martin, R. Erik (see Proffit, C. Edward)	
Matsui, Masafumi. Geographic variation in toads of the <i>Bufo bufo</i> complex from the Far East, with a description of a new subspecies	561
Matsui, Masafumi. Three new species of <i>Amolops</i> from Borneo (Amphibia, Anura, Ranidae)	623
Matthews, William J. Fish faunal structure in an Ozark stream: stability, persistence and a catastrophic flood	388
McClanahan, Lon L., C. Robert Feldmeth, Jeff Jones and David L. Soltz. Energetics, salinity and temperature tolerance in the Mohave tui chub, <i>Gila bicolor mohavensis</i>	45
McKaye, Kenneth R. and Jay R. Stauffer, Jr. Description of a gold cichlid (Teleostei: Cichlidae) from Lake Malawi, Africa	870
McKelvie, D. Scott (see Methven, David A.)	
Methven, David A. and D. Scott McKelvie. Distribution of <i>Phycis chesteri</i> (Pisces: Gadidae) on the Grand Bank and Labrador shelf	886
Miller, Robert Rush (see Chernoff, Barry)	
Minckley, C. O. (see Minckley, W. L.)	
Minckley, W. L. and C. O. Minckley. <i>Cyprinodon pachycephalus</i> , a new species of pupfish (Cyprinodontidae) from the Chihuahuan Desert of northern México	184
Miyamoto, Michael M. Phylogenetic relationships and systematics of the <i>Eleutherodactylus fitzingeri</i> group (Anura, Leptodactylidae)	503
Miyamoto, Michael M., Marc P. Hayes and Michele R. Tennant. Biochemical and morphological variation in Floridian populations of the bark anole ( <i>Anolis distichus</i> )	76
Moler, Paul E. (see Anderson, Karen)	
Morreale, S. J. (see Scribner, K. T.)	
Moring, John R. and Matthew E. Moring. A late leptocephalus stage of a conger eel, <i>Conger oceanicus</i> , found in a tidepool	222
Moring, Matthew E. (see Moring, John R.)	
Morman, R. H. (see Jacobson, L. D.)	
Moyer, Jack T. Longevity of the anemonefish <i>Amphiprion clarkii</i> at Miyake-jima, Japan, with notes on four other species	135
Muldoon, Kathryn A. (see Proffit, C. Edward)	
Munger, James C. Rate of death due to predation for two species of horned lizard, <i>Phrynosoma cornutum</i> and <i>P. modestum</i>	820
Natanson, Lisa J. and Gregor M. Cailliet. Reproduction and development of the Pacific angel shark, <i>Squatina californica</i> , off Santa Barbara, California	987

NELSON, GARETH. Identity of the anchovy <i>Engraulis clarki</i> with notes on the species-groups of <i>Anchoa</i>	891
NIELSEN, JØRGEN G. <i>Leptobrotula breviventralis</i> , a new bathyal fish genus and species from the Indo-west Pacific (Ophidiiformes, Ophidiidae)	166
NOONAN, WILLIAM R. (see Kaeding, Lynn R.)	
NORELL, MARK ALLEN. Late Pleistocene lizards from Kokoweef Cave, San Bernardino County, California	244
PAGE, LAWRENCE M. (see Retzer, Michael E.)	
PARENTI, LYNNE R. Homology of pelvic fin structures in female phallostethid fishes (Atherinomorpha, Phallostethidae)	305
PARKER, NICK C. (see Thompson, Kenneth W.)	
PATCHELL, FREDERICK C. AND RICHARD SHINE. Food habits and reproductive biology of the Australian legless lizards (Pygopodidae)	30
PEERY, BRUCE D. (see Proffit, C. Edward)	
PENNYPACKER, KEITH R. (see Fine, Michael L.)	
PIETSCH, THEODORE W. The original manuscript sources for the Histoire Naturelle Des Poissons, 1828—1849: keys to understanding the fishes described by Cuvier and Valenciennes	216
PIETSCH, THEODORE W. Systematics and distribution of bathypelagic anglerfishes of the family Ceratiidae (Order: Lophiiformes)	479
PORTER, WARREN P. (see Christian, Keith A.)	
PROFFIT, C. EDWARD, R. ERIK MARTIN, ROBERT G. ERNST, BARBARA J. GRAUNKE, SARA E. LECROY, KATHRYN A. MULDOON, BRUCE D. PEERY, J. ROSS WILCOX AND NANCY WILLIAMS-WALLS. Effects of power plant construction and operation on the nesting of the loggerhead sea turtle ( <i>Caretta caretta</i> ): 1971–84	813
RANDALL, JOHN E. (see Fraser, Thomas H.)	
RETZER, MICHAEL E., LAWRENCE M. PAGE AND DAVID L. SWOFFORD. Variation and systematics of <i>Etheostoma whipplei</i> , the redfin darter (Pisces: Percidae)	631
RIVAS, LUIS R. Systematic review of the perciform fishes of the genus <i>Centropomus</i>	579
ROMANI, ROMANO (see Torticelli, Patrizia)	
ROSENBLATT, RICHARD H. AND ROBIN S. WAPLES. A genetic comparison of allopatric populations of shore fish species from the eastern and central Pacific Ocean: dispersal or vicariance?	275
ROSS, STEPHEN T. Resource partitioning in fish assemblages: a review of field studies	352
SAINSBURY, K. J. (see Thresher, R. E.)	
SCHAEFER, KURT M. Lethal temperatures and the effect of temperature change on volitional swimming speeds of chub mackerel, <i>Scomber japonicus</i>	39
SCHAEFER, SCOTT A. AND TED M. CAVENDER. Geographic variation and subspecific status of <i>Notropis spilopterus</i> (Pisces: Cyprinidae)	122
SCHRAEDER, PATRICIA A. (see Kaeding, Lynn R.)	
SCHUETT, GORDON W. AND JAMES C. GILLINGHAM. Sperm storage and multiple paternity in the copperhead, <i>Agkistrodon contortrix</i>	807
SCHULTZ, ERIC T. <i>Pterois volitans</i> and <i>Pterois miles</i> : two valid species	686
SCHWARTZ, ALBERT (see Henderson, Robert W.)	
SCRIBNER, K. T., J. E. EVANS, S. J. MORREALE, M. H. SMITH AND J. W. GIBBONS. Genetic divergence among populations of the yellow-bellied slider turtle ( <i>Pseudemys scripta</i> ) separated by aquatic and terrestrial habitats	691
SEIDEL, MICHAEL E., JOHN B. IVERSON AND M. DALE ADKINS. Biochemical comparisons and phylogenetic relationships in the family Kinosternidae (Testudines)	285
SELCEK, KYLE W. Life history of a successful colonizer: the Mediterranean gecko, <i>Hemidactylus turcicus</i> , in southern Texas	956
SHINE, RICHARD (see Patchell, Frederick C.)	
SHINE, RICHARD. Predation upon filesnakes ( <i>Acrochordus arafurae</i> ) by aboriginal hunters: selectivity with respect to body size, sex and reproductive condition	238
SHINE, RICHARD. Ecology of a low-energy specialist: food habits and reproductive biology of the Arafura filesnake ( <i>Acrochordidae</i> )	424
SHINE, RICHARD. Evolutionary advantages of limlessness: evidence from the pygopodid lizards	525
SHUTTLESWORTH, GLADIS A. (see Ford, Neil B.)	
SMITH, MICHAEL H. (see Scribner, K. T.)	
SMITH, MICHAEL H. (see Kennedy, Phyllis K.)	
SMITH, P. J. Low genetic variation in sharks (Chondrichthyes)	202
SMITS, ALLAN W., JOHN WARD AND HARVEY LILLYWHITE. Effects of hyperkalemia on thermoregulatory and feeding behaviors of the lizard, <i>Sauromalus hispidus</i>	518

SNELSON, FRANKLIN F., JR., JEFFREY D. WETHERINGTON AND HELEN L. LARGE. The relationship between interbrood interval and yolk loading in a generalized poeciliid fish, <i>Poecilia latipinna</i> .....	295
SNYDER, DAVID B. (see Taylor, Jeffrey N.)	
SOLTZ, DAVID L. (see McClanahan, Lon L.)	
SOUTHERLAND, MARK T. Behavioral niche expansion in <i>Desmognathus fuscus</i> (Amphibia: Caudata: Plethodontidae) .....	235
SOUTHERLAND, MARK T. The effects of variation in streamside habitats on the composition of mountain salamander communities .....	731
SPRIESTERSBACH, KARLA K. (see Hutchison, Victor H.)	
STARNES, WAYNE C. (see Etnier, David A.)	
STAUFFER, JAY R., JR. (see McKaye, Kenneth R.)	
STAUFFER, JAY R., JR. (see Boltz, J. M.)	
STEWART, DONALD J. (see Leslie, Andrew J., Jr.)	
STEWART, DONALD J. Revision of <i>Pimelodina</i> and description of a new genus and species from the Peruvian Amazon (Pisces: Pimelodidae) .....	653
STEWART, MARGARET M. (see Townsend, Daniel S.)	
SWOFFORD, DAVID L. (see Retzer, Michael E.)	
TABER, CHARLES A. (see James, Paul W.)	
TAVE, DOUGLAS. A quantitative genetic analysis of 19 phenotypes in <i>Tilapia nilotica</i> .....	672
TAYLOR, JANET A. Seasonal energy storage in the Australian lizard, <i>Ctenotus taeniolatus</i> .....	445
TAYLOR, JEFFREY N., DAVID B. SNYDER AND WALTER R. COURTENAY, JR. Hybridization between two introduced, substrate-spawning tilapias (Pisces: Cichlidae) in Florida .....	903
TENNANT, MICHELE R. (see Miyamoto, Michael M.)	
THOMPSON, KENNETH W., LUTHER A. KNIGHT, JR. AND NICK C. PARKER. Color-coded fluorescent plastic chips for marking small fishes .....	544
THRESHER, R. E., K. J. SAINSBURY, J. S. GUNN AND A. W. WHITELAW. Life history strategies and recent changes in population structure in the lizardfish genus <i>Saurida</i> , on the Australian northwest shelf .....	876
TINKLE, DONALD W. AND ARTHUR E. DUNHAM. Comparative life histories of two syntopic sceloporine lizards .....	1
TODD, THOMAS N. Occurrence of white bass-white perch hybrids in Lake Erie .....	196
TORBLAA, R. L. (see Jacobson, L. D.)	
TORRICELLI, PATRIZIA AND ROMANO ROMANI. Sound production in the Italian freshwater goby, <i>Padogobius martensi</i> .....	213
TOWNSEND, DANIEL S. AND MARGARET M. STEWART. The effect of temperature on direct development in a terrestrial-breeding, neotropical frog .....	520
TRACY, C. RICHARD (see Christian, Keith A.)	
TRAVIS, JOSEPH (see Fair, James A.)	
TURNER, FREDERICK B. (see Betty, Kristin H.)	
ULTSCH, GORDON R. AND JOHN F. ANDERSON. The respiratory microenvironment within the burrows of gopher tortoises ( <i>Gopherus polyphemus</i> ) .....	787
UWA, HIROSHI AND WICHIAN MAGTOON. Description and karyotype of a new ricefish, <i>Oryzias mekongensis</i> , from Thailand .....	473
VAN DRAGT, RANDALL G. (see Kats, Lee B.)	
VITT, LAURIE J. Reproductive tactics of sympatric gekkonid lizards with a comment on the evolutionary and ecological consequences of invariant clutch size .....	773
VITTURI, R., P. CARBONE, M. MACALUSCO AND E. CATALANO. Karyotypic variability in the gadid fish, <i>Gaidropsarus mediterraneus</i> (Linnaeus, 1758) .....	826
VORIS, HAROLD K. (see Inger, Robert F.)	
WAGNER, WILLIAM E., JR. Tadpoles and pollen: observations on the feeding behavior of <i>Hyla regilla</i> larvae .....	802
WALKER, PAUL (see Inger, Robert F.)	
WAPLES, ROBIN S. (see Rosenblatt, Richard H.)	
WARD, JOHN (see Smits, Allan W.)	
WERMAN, STEVEN D. Phylogenetic relationships of the true viper, <i>Eristocophis mcmahoni</i> , based on parsimony analysis of allozyme characters .....	1014
WEST, ROBIN L. (see Goldberg, Stephen R.)	
WETHERINGTON, JEFFREY D. (see Snelson, Franklin F., Jr.)	
WHITELAW, A. W. (see Thresher, R. E.)	
WHITMAN, PAMELA A., JOSEPH A. MARSHALL AND E. C. KELLER, JR. Tonic immobility in the smooth dogfish shark, <i>Mustelus canis</i> (Pisces, Carcharhinidae) .....	829
WILCOX, J. ROSS (see Proffit, C. Edward)	

WILEY, JOHN E. AND ALVIN L. BRASWELL. A triploid male <i>Rana palustris</i> .....	531
WILLIAMS-WALLS, NANCY (see Proffit, C. Edward)	
WOODMAN, DAVID A. Microgeographic variation in <i>Rhinichthys cataractae</i> (Cyprinidae) in western Nebraska .....	219
YASUTAKE, WILLIAM T. (see Goldberg, Stephen R.)	
ZALISKO, EDWARD J. (see Beneski, John T., Jr.)	
ZAPPALORTI, ROBERT T. (see Burger, Joanna)	
ZIMMERMAN, EARL G. (see Kennedy, Phyllis K.)	